

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 March 2003 (27.03.2003)

PCT

(10) International Publication Number
WO 03/024928 A2

(51) International Patent Classification⁷: C07D 207/14,
211/56, 405/12, 205/04, A61K 31/40, 31/445, 31/4523,
A61P 1/04, 3/04, 3/10, 25/28, 37/08

(21) International Application Number: PCT/DK02/00593

(22) International Filing Date:
11 September 2002 (11.09.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PA 2001 01344 14 September 2001 (14.09.2001) DK
PA 2002 00750 16 May 2002 (16.05.2002) DK

(71) Applicants: NOVO NORDISK A/S [DK/DK]; Novo
Allé, DK-2880 Bagsværd (DK). BOEHRINGER IN-
GELHEIM INTERNATIONAL GMBH [DE/DE];
55216 Ingelheim am Rhein (DE).

(72) Inventors: DÖRWALD, Florencio, Zaragoza; Højager-
parken 30, 1, DK-2750 Ballerup (DK). HOHLWEG, Rolf,
Nybovej 6, DK-3490 Kvistgaard (DK).

(74) Common Representative: NOVO NORDISK A/S; Cor-
porate Patents, Novo Allé, DK-2880 Bagsværd (DK).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC,
VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.



WO 03/024928 A2

(54) Title: NOVEL AMINOAZETIDINE, -PYRROLIDINE AND -PIPERIDINE DERIVATIVES

(57) Abstract: Novel aminoazetidine, -pyrrolidine, and -piperidine derivatives, use of these compounds as pharmaceutical compositions, pharmaceutical compositions comprising the compounds, and a method of treatment employing these compounds and compositions. The compounds show a high and selective binding affinity to the histamine H3 receptor indicating histamine H3 receptor antagonistic, inverse agonistic or agonistic activity. As a result, the compounds are useful for the treatment of diseases and disorders related to the histamine H3 receptor.

NOVEL AMINOAZETIDINE, -PYRROLIDINE AND -PIPERIDINE DERIVATIVES

FIELD OF THE INVENTION

The present invention relates to novel aminoazetidine, -pyrrolidine and -piperidine derivatives, to the use of these compounds as pharmaceutical compositions, to pharmaceutical compositions comprising the compounds, and to a method of treatment employing these compounds and compositions. The present compounds show a high and selective binding affinity to the histamine H3 receptor indicating histamine H3 receptor antagonistic, inverse-agonistic or agonistic activity. As a result, the compounds are useful for the treatment of diseases and disorders related to the histamine H3 receptor.

10 BACKGROUND OF THE INVENTION

The existence of the histamine H3 receptor has been known for several years and the receptor is of current interest for the development of new medicaments (see eg Stark et al., *Drugs Fut.* **1996**, *21*, 507-520; Leurs et al., *Progress in Drug Research* **1995**, *45*, 107-165). Recently, the human histamine H3 receptor has been cloned, cf Lovenberg et al., *Molecular Pharmacology*, **June 1999**, *55*, 1101-1107. The histamine H3 receptor is a presynaptic autoreceptor located both in the central and the peripheral nervous system, the skin and in organs such as the lung, the intestine, probably the spleen and the gastrointestinal tract. Recent evidence suggests that the H3 receptor shows intrinsic, constitutive activity, *in vitro* as well as *in vivo* (ie it is active in the absence of an agonist; see eg Morisset et al., *Nature* **2000**, *408*, 860-864). Compounds acting as inverse agonists can inhibit this activity. The histamine H3 receptor has been demonstrated to regulate the release of histamine and also of other neurotransmitters such as serotonin and acetylcholine. A histamine H3 receptor antagonist or inverse agonist would therefore be expected to increase the release of these neurotransmitters in the brain. A histamine H3 receptor agonist, on the contrary, leads to an inhibition of the biosynthesis of histamine and an inhibition of the release of histamine and also of other neurotransmitters such as serotonin and acetylcholine. These findings suggest that histamine H3 receptor agonists, inverse agonists and antagonists could be important mediators of neuronal activity. Accordingly, the histamine H3 receptor is an important target for new therapeutics.

Compounds similar to the compounds of the present invention have previously been disclosed, cf US 3,963,745, US 3,966,957, DE 25 58 348, US 6,090,802, WO 97/23483, WO 97/09308, JP 07048375, EP 518 558, *Acta Pharm Suec. Suppl.* **1**, 200-218, **1983**, *Life Sci.*, **32**(25), 2877-2886, **1983**, US 4,109,005, BE 861790, *Aviat. Space Environ. Med.*, **48**(9),

867-871, 1977, US 4,049,821, US 4,039,678, *Therapie* 1977, 32(1), 121-132, US 4,002,757 DE 19 64 516, NL 6500326, US 3,787,419, Irikura, *Yakugaku Zasshi* 1962, 82, 356, DE 24 61 802, *Eur. J. Chem. Chim. Ther.* 1982, 17, 437, *J. Med. Chem.* 1992, 35, 4334, *Chem. Pharm. Bull.* 1995, 43, 947, *Arzneim. Forsch.* 1999, 49, 96, US 5,571,832, *NIDA Res.*

- 5 *Monogr.* 1993, 140 (Problems of Drug Dependence, volume 1), 179-195. However, these references neither disclose nor suggest that these compounds may have a histamine H3 receptor antagonistic or agonistic activity.

Several publications disclose the preparation and use of histamine H3 agonists and antagonists. Most of these are imidazole derivatives (see eg Stark et al., *Drugs of the Future* 10 1996, 21, 507-520; Tozer, Kalindjian, *Expert Opinion on Therapeutic Patents* 2000, 10, 1045-1055). However, recently some imidazole-free ligands of the histamine H3 receptor have been described (see eg Walczynski et al., *Arch. Pharm. Pharm. Med. Chem.* 1999, 332, 389-398; Linney et al., *J. Med. Chem.* 2000, 43, 2362-2370; Ganellin et al., *Arch. Pharm. Pharm. Med. Chem.* 1998, 331, 395-404; Walczynski et al., *Il Farmaco* 1999, 54, 15 684-694; WO 99/42458, EP 0 978 512, WO 97/17345, US 6,316,475, WO 01/66534, WO 01/74810, WO 01/44191, WO 01/74815, WO 01/74773, WO 01/74813, WO 01/74814 and WO 02/12190.


However, these compounds differ structurally from the present compounds.

In view of the art's interest in histamine H3 receptor agonists, inverse agonists and 20 antagonists, novel compounds which interact with the histamine H3 receptor would be a highly desirable contribution to the art. The present invention provides such a contribution to the art being based on the finding that a novel class of aminoazetidine, -pyrrolidine and -piperidine derivatives has a high and specific affinity to and potency at the histamine H3 receptor.

25 Due to their interaction with the histamine H3 receptor, the present compounds are useful in the treatment of a wide range of conditions and disorders in which an interaction with the histamine H3 receptor is beneficial. Thus, the compounds may find use eg in the treatment of diseases of the central nervous system, the peripheral nervous system, the cardiovascular system, the pulmonary system, the gastrointestinal system and the endocrino- 30 logical system.

SUMMARY OF THE INVENTION

The invention relates to a compound of the general formula (I):



5 wherein

R^1 represents

- hydrogen,
- 10 • C_{1-8} -alkyl, C_{3-8} -alkenyl or C_{3-8} -alkynyl,
which may optionally be substituted with one or more halogen atoms,
- C_{3-7} -cycloalkyl, C_{3-7} -cycloalkenyl, C_{4-8} -bicycloalkyl, C_{3-7} -cycloalkyl- C_{1-3} -alkyl or
15 C_{3-7} -cycloalkenyl- C_{1-3} -alkyl,
which may optionally be substituted at any position with one or more halogen atoms,

R^2 represents C_{1-6} -alkyl,

20 f is 0, 1 or 2,

X represents $-(CH_2)_m-(Z)_n-(CH_2)_o-$,

m and o independently are 0, 1, 2, 3 or 4,

25

n is 0 or 1,

Z is $-O-$, $-NH-$, $-N(CH_3)-$, $-C(=O)-$, $-CH(OH)-$, $-CH(O-C_{1-6}-alkyl)-$, $-C(=N-OH)-$, $-S-$, $-S(=O)-$,
 $-S(=O)_2-$, $-CH=CH-$ or $-C\equiv C-$,

30

Y is

(a) aryl or heteroaryl, which may optionally be substituted with one or more substituents selected from

- 5 ▪ halogen, nitro, cyano, hydroxy, C₁₋₆-alkanoyl, C₁₋₆-alkylthio, C₁₋₆-alkylsulfonyl, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₃₋₈-cycloalkyl, trifluoromethyl, trifluoromethoxy, -NR³R⁴ and -O(C=O)NR³R⁴, or wherein two substituents in adjacent positions form a radical -O-(CH₂)₁₋₃-O-, wherein R³ and R⁴ independently are hydrogen, C₁₋₆-alkyl, C₃₋₈-cycloalkyl,

10 C₁₋₆-alkanoyl or aryl, or R³ and R⁴ together with the nitrogen atom to which they are attached form a 4 to 7 membered, saturated or unsaturated azetidyl, pyrrolidinyl, piperidyl or azepanyl ring,

 15 ▪ aryl, aryloxy, aryl-C₁₋₆-alkyl and aryl-C₁₋₆-alkoxy, wherein the ring moieties optionally may be substituted with one or more substituents selected from
 - halogen, nitro, cyano, hydroxy, C₁₋₆-alkanoyl, C₁₋₆-alkylthio, C₁₋₆-alkylsulfonyl, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₃₋₈-cycloalkyl, trifluoromethyl, trifluoromethoxy, -NR⁵R⁶ and -O(C=O)NR⁵R⁶, or wherein two substituents in adjacent positions form a radical -O-(CH₂)₁₋₃-O-,

20 wherein R⁵ and R⁶ independently are hydrogen, C₁₋₆-alkyl, C₃₋₈-cycloalkyl, C₁₋₆-alkanoyl or aryl, or R⁵ and R⁶ together with the nitrogen atom to which they are attached form a 4 to 7 membered, saturated or unsaturated azetidyl, pyrrolidinyl, piperidyl or azepanyl ring,

25 (b) C₃₋₈-cycloalkyl or C₅₋₈-cycloalkenyl, which may optionally be substituted with one or more substituents selected from
 - C₁₋₆-alkyl, C₁₋₆-alkoxy, C₁₋₆-alkylthio, cyano, trifluoromethyl, trifluoromethoxy and halogen,

30 aryl and aryloxy, wherein the ring moieties optionally may be substituted with one or more substituents selected from
 - halogen, nitro, cyano, hydroxy, C₁₋₆-alkanoyl, C₁₋₆-alkylthio, C₁₋₆-alkylsulfonyl, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₃₋₈-cycloalkyl, trifluoromethyl, trifluoromethoxy, -NR⁷R⁸ and -O(C=O)NR⁷R⁸, or wherein two substituents in adjacent positions form a radical -O-(CH₂)₁₋₃-O-,

35

wherein R^7 and R^8 independently are hydrogen, C_{1-6} -alkyl, C_{3-8} -cycloalkyl, C_{1-6} -alkanoyl or aryl, or R^7 and R^8 together with the nitrogen atom to which they are attached form a 4 to 7 membered, saturated or unsaturated azetid-
inyl, pyrrolidinyl, piperidyl or azepanyl ring,

5

as well as any diastereomer or enantiomer or tautomeric form thereof including mixtures of these or a pharmaceutically acceptable salt thereof,



with the proviso that,

10


when m, n, and o are 0, R^1 must not be cyclopentyl, cyclohexyl, ethyl, or methyl,




and with the proviso that the compound must not be

15




20





or



5

DEFINITIONS

In the structural formulae given herein and throughout the present specification, the following terms have the indicated meaning:

10 The term "halogen" means F, Cl, Br or I.

The term "C₁₋₆-alkyl" as used herein represents a saturated, branched or straight hydrocarbon group having from 1 to 6 carbon atoms. Typical C₁₋₆-alkyl groups include, but are not limited to, methyl, ethyl, n-propyl, isopropyl, butyl, isobutyl, *sec*-butyl, *tert*-butyl, pentyl, hexyl and the like.

15 In the same way the terms "C₁₋₃-alkyl" and "C₁₋₈-alkyl" as used herein represent saturated, branched or straight hydrocarbon groups having from 1 to 3 carbon atoms and from 1 to 8 carbon atoms, respectively.

The term "C₃₋₈-alkenyl" as used herein represents a branched or straight hydrocarbon group having from 3 to 8 carbon atoms and at least one double bond. Typical C₃₋₈-alkenyl groups include, but are not limited to, 1-propenyl, 2-propenyl, isopropenyl, 1,3-butadienyl, 1-butenyl, 2-butenyl, 1-pentenyl, 2-pentenyl, 1-hexenyl, 2-hexenyl, 2-

20

heptenyl, 2-octenyl, 1-ethylprop-2-enyl, 1,1-(dimethyl)prop-2-enyl, 1-ethylbut-3-enyl, 1,1-(dimethyl)but-2-enyl, 1,1-(dimethyl)pent-3-enyl, 1-ethylpent-2-enyl, 1,1-(dimethyl)pent-3-enyl, 1,1-(dimethyl)hex-3-enyl, 1-ethylhex-4-enyl and the like.

The term "C₃₋₈-alkynyl" as used herein represents a branched or straight hydrocarbon group having from 3 to 8 carbon atoms and at least one triple bond. Typical C₃₋₈-alkynyl groups include, but are not limited to, 1-propynyl, 2-propynyl, isopropynyl, 1,3-butadiynyl, 1-butynyl, 2-butynyl, 1-pentynyl, 2-pentynyl, 1-hexynyl, 2-hexynyl, 2-heptynyl, 2-octynyl, 1-ethylprop-2-ynyl, 1,1-(dimethyl)prop-2-ynyl, 1-ethylbut-3-ynyl, 1,1-(dimethyl)but-2-ynyl, 1,1-(dimethyl)pent-3-ynyl, 1-ethylpent-2-ynyl, 1,1-(dimethyl)pent-3-ynyl, 1,1-(dimethyl)hex-3-ynyl, 1-ethylhex-4-ynyl and the like.

The term "C₁₋₆-alkoxy" as used herein refers to the radical -O-C₁₋₆-alkyl, wherein C₁₋₆-alkyl is as defined above. Representative examples are methoxy, ethoxy, n-propoxy, isopropoxy, butoxy, *sec*-butoxy, *tert*-butoxy, pentoxy, isopentoxy, hexoxy, isohexoxy and the like.

The term "C₁₋₆-alkanoyl" as used herein refers to the radical -C(=O)H or -C(=O)C₁₋₅-alkyl, wherein C₁₋₅-alkyl represents a saturated, branched or straight hydrocarbon groups having from 1 to 5 carbon atoms. Representative examples are acetyl, propionyl, butanoyl, pentanoyl, hexanoyl, and the like.

The term "C₁₋₆-alkylthio" as used herein refers to the radical -S-C₁₋₆-alkyl, wherein C₁₋₆-alkyl represents a saturated, branched or straight hydrocarbon groups having from 1 to 6 carbon atoms as defined above. Representative examples are methylthio, ethylthio, propylthio, butylthio, pentylthio, hexylthio, and the like.

The term "C₁₋₆-alkylsulfonyl" as used herein refers to the radical -S(=O)₂-C₁₋₆-alkyl, wherein C₁₋₆-alkyl represents a saturated, branched or straight hydrocarbon groups having from 1 to 6 carbon atoms as defined above. Representative examples are methylsulfonyl, ethylsulfonyl, propylsulfonyl, butylsulfonyl, pentylsulfonyl, hexylsulfonyl, and the like.

The term "C₃₋₇-cycloalkyl" as used herein represents a saturated, monocyclic, carbocyclic group having from 3 to 7 carbon atoms. Representative examples are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and the like.

The term "C₃₋₇-cycloalkenyl" as used herein represents a monocyclic, carbocyclic, non-aromatic group having from 3 to 7 carbon atoms and at least one double bond. Representative examples are cyclopropenyl, cyclobutenyl, cyclopentenyl and the like.

In the same way the term "C₅₋₈-cycloalkenyl" as used herein represent a monocyclic, carbocyclic, non-aromatic group having from 5 to 8 carbon atoms and at least one double bond.

The term "C₄₋₈-bicycloalkyl" as used herein represents a saturated, bicyclic, carbocyclic group having from 4 to 8 carbon atoms. Representative examples are bicyclo[3.1.0]hexyl, bicyclo[4.1.0]heptyl, bicyclo[3.1.1]heptyl, bicyclo[2.2.1]heptyl and the like.


5 The term "aryl" as used herein is intended to include carbocyclic aromatic ring systems such as phenyl, biphenyl, naphthyl, anthracenyl, phenanthrenyl, fluorenyl, indenyl, pentalenyl, azulenyl and the like. Aryl is also intended to include the partially hydrogenated derivatives of the carbocyclic systems enumerated above. Non-limiting examples of such partially hydrogenated derivatives are 1,2,3,4-tetrahydronaphthyl, 1,4-dihydronaphthyl and the like.

10 The term "aryloxy" as used herein refers to the radical -O-aryl, wherein aryl is as defined above. Non-limiting examples are phenoxy, naphthoxy, anthracenyloxy, phenantrenyloxy, fluorenyloxy, indenyloxy and the like.

The term "heteroaryl" as used herein is intended to include heterocyclic aromatic ring systems containing one or more heteroatoms selected from nitrogen, oxygen and sulfur such as furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, isoxazolyl, isothiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, pyran, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, 1,2,3-triazinyl, 1,2,4-triazinyl, 1,3,5-triazinyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, tetrazolyl, thiazidinyl, indolyl, isoindolyl, benzofuryl, benzothienyl, indazolyl, benzimidazolyl, benzothiazolyl, benzoisothiazolyl, benzoxazolyl, benzisoxazolyl, purinyl, quinazolinyl, quinolizinyll, isoquinolinyll, quinoxalinyll, naphthyridinyll, pteridinyll, carbazolyll, azepinyll, diazepinyll, acridinyll and the like. Heteroaryl is also intended to include the partially hydrogenated derivatives of the heterocyclic systems enumerated above. Non-limiting examples of such partially hydrogenated derivatives are 2,3-dihydrobenzofuranyl, pyrrolinyll, pyrazolinyll, indanyll, indolinyll, oxazolidinyll, oxazolinyll, oxazepinyll and the like.

25 Certain of the above defined terms may occur more than once in the structural formulae, and upon such occurrence each term shall be defined independently of the other.

"Aryl-C₁₋₆-alkyl", "aryl-C₁₋₆-alkoxy" etc. mean C₁₋₆-alkyl or C₁₋₆-alkoxy as defined above, substituted by aryl as defined above, for example:



30 The term "optionally substituted" as used herein means that the groups in question are either unsubstituted or substituted with one or more of the substituents specified. When

the groups in question are substituted with more than one substituent the substituents may be the same or different.

The term "treatment" as used herein means the management and care of a patient for the purpose of combating a disease, disorder or condition. The term is intended to include the delaying of the progression of the disease, disorder or condition, the alleviation or relief of symptoms and complications, and/or the cure or elimination of the disease, disorder or condition. The patient to be treated is preferably a mammal, in particular a human being.

DESCRIPTION OF THE INVENTION

In one embodiment, Y is

(a) aryl or heteroaryl, which may optionally be substituted with one or more substituents selected from

- halogen, nitro, cyano, hydroxy, C₁₋₆-alkanoyl, C₁₋₆-alkylthio, C₁₋₆-alkylsulfonyl, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₃₋₈-cycloalkyl, trifluoromethyl, trifluoromethoxy, -NR³R⁴ and -O(C=O)NR³R⁴, or wherein two substituents in adjacent positions form a radical -O-(CH₂)₁₋₃-O-,
wherein R³ and R⁴ independently are hydrogen, C₁₋₆-alkyl, C₃₋₈-cycloalkyl, C₁₋₆-alkanoyl or aryl, or R³ and R⁴ together with the nitrogen atom to which they are attached form a 4 to 7 membered, saturated or unsaturated azetidyl, pyrrolidinyl, piperidyl or azepanyl ring,
- aryl, aryl-C₁₋₆-alkyl and aryl-C₁₋₆-alkoxy, wherein the ring moieties optionally may be substituted with one or more substituents selected from
 - halogen, nitro, cyano, hydroxy, C₁₋₆-alkanoyl, C₁₋₆-alkylthio, C₁₋₆-alkylsulfonyl, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₃₋₈-cycloalkyl, trifluoromethyl, trifluoromethoxy, -NR⁵R⁶ and -O(C=O)NR⁵R⁶, or wherein two substituents in adjacent positions form a radical -O-(CH₂)₁₋₃-O-,
wherein R⁵ and R⁶ independently are hydrogen, C₁₋₆-alkyl, C₃₋₈-cycloalkyl, C₁₋₆-alkanoyl or aryl, or R⁵ and R⁶ together with the nitrogen atom to which they are attached form a 4 to 7 membered, saturated or unsaturated azetidyl, pyrrolidinyl, piperidyl or azepanyl ring,

(b) C₃₋₈-cycloalkyl or C₅₋₈-cycloalkenyl, which may optionally be substituted with one or more substituents selected from

- C₁₋₆-alkyl, C₁₋₆-alkoxy, C₁₋₆-alkylthio, cyano, trifluoromethyl, trifluoromethoxy and halogen,
- aryl and aryloxy, wherein the ring moieties optionally may be substituted with one or more substituents selected from
 - halogen, nitro, cyano, hydroxy, C₁₋₆-alkanoyl, C₁₋₆-alkylthio, C₁₋₆-alkylsulfonyl, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₃₋₈-cycloalkyl, trifluoromethyl, trifluoromethoxy, -NR⁷R⁸ and -O(C=O)NR⁷R⁸, or wherein two substituents in adjacent positions form a radical -O-(CH₂)₁₋₃-O-, wherein R⁷ and R⁸ independently are hydrogen, C₁₋₆-alkyl, C₃₋₈-cycloalkyl, C₁₋₆-alkanoyl or aryl, or R⁷ and R⁸ together with the nitrogen atom to which they are attached form a 4 to 7 membered, saturated or unsaturated azetidiny, pyrrolidinyl, piperidyl or azepanyl ring,

In another embodiment, f is 1.

In another embodiment, R¹ is C₁₋₈-alkyl, C₃₋₇-cycloalkyl-C₁₋₃-alkyl or C₃₋₇-cycloalkyl, which may optionally be substituted with one or more halogen atoms,

In still another embodiment, R¹ is C₁₋₈-alkyl, C₃₋₇-cycloalkyl-C₁₋₃-alkyl or C₃₋₇-cycloalkyl, such as 1-ethylpropyl, 2-propyl, cyclobutyl or cyclopentyl.

In yet another embodiment, R² is methyl.

In a further embodiment, X is -(CH₂)_m-C(=O)-CH₂-, -(CH₂)_m-, a valence bond or -(CH₂)_m-CH=CH-(CH₂)_o-, wherein m and o are as defined for formula (I).

In yet a further embodiment, X is -(CH₂)_m-C(=O)-, -(CH₂)_m-, a valence bond or -CH=CH-, wherein m is 1, 2, 3 or 4, such as -(CH₂)₂-C(=O)-, -CH₂-, -(CH₂)₂-, a valence bond or -CH=CH-.

In still a further embodiment, X is -(CH₂)_m-C(=O)-, -(CH₂)_m- or -CH=CH-, wherein m is 1, 2, 3 or 4, such as -(CH₂)₂-C(=O)-, -CH₂-, -(CH₂)₂- or -CH=CH-.

In another embodiment, Y is aryl, such as phenyl, which may optionally be substituted as defined for formula (I).

5 In still another embodiment, Y is phenyl, which may optionally be substituted with one or two of

- C₁₋₆-alkoxy, -NR³R⁴, halogen, trifluoromethyl or hydroxy, wherein R³ and R⁴ are as defined for formula (I),

10

- two substituents in adjacent positions forming a radical -O-(CH₂)₁₋₃-O-, or
- phenyl or phenoxy, which may optionally be substituted as defined for formula (I).


15 In yet another embodiment, Y is phenyl, which may optionally be substituted with one or two of

- C₁₋₆-alkoxy, -NR³R⁴, halogen, trifluoromethyl or hydroxy, wherein R³ and R⁴ independently are hydrogen or C₁₋₆-alkyl,

20


- two substituents in adjacent positions forming a radical -O-CH₂-O-, or
- phenyl or phenoxy.

25 In another embodiment the invention relates to a compound of the general formula (I₁):




wherein R¹, R² and Y are as defined for formula (I) or any of the above embodiments.

In still another embodiment the invention relates to a compound of the general formula (I₂):



In yet another embodiment the invention relates to a compound of the general formula (I₃):

5



In still a further embodiment, the compound is selected from

- 4-(3,4-dimethoxyphenyl)-N-[1-(1-ethylpropyl)pyrrolidin-3-yl]-N-methyl-4-oxobutyramide,
- 2-(4-dimethylaminophenyl)-N-[1-(1-ethylpropyl)pyrrolidin-3-yl]-N-methylacetamide,
- 10 2-biphenyl-4-yl-N-[1-(1-ethylpropyl)pyrrolidin-3-yl]-N-methylacetamide,
- 4-(4-chlorophenyl)-N-[1-(1-ethylpropyl)pyrrolidin-3-yl]-N-methyl-4-oxobutyramide,
- N-[1-(1-ethylpropyl)pyrrolidin-3-yl]-N-methyl-3-(4-trifluoromethylphenyl)propionamide,
- N-[1-(1-ethylpropyl)pyrrolidin-3-yl]-N-methyl-3-phenoxybenzamide,
- 4-(3,4-dimethoxyphenyl)-N-[1-(1-ethylpropyl)pyrrolidin-3-yl]-4-oxobutyramide,
- 15 N-[1-(1-ethylpropyl)pyrrolidin-3-yl]-2-(3-fluoro-4-hydroxyphenyl)-N-methylacetamide,
- 2-biphenyl-4-yl-N-[1-(1-ethylpropyl)pyrrolidin-3-yl]-N-methylacetamide,
- 2-biphenyl-4-yl-N-(1-cyclopentylpyrrolidin-3-yl)-N-methylacetamide,
- 3-benzo[1,3]dioxol-5-yl-N-[1-(1-ethylpropyl)pyrrolidin-3-yl]-N-methylacrylamide,
- 3-(3-chlorophenyl)-N-[1-(1-ethylpropyl)pyrrolidin-3-yl]-N-methylacrylamide,
- 20 2-biphenyl-4-yl-N-(1-isopropylpyrrolidin-3-yl)-N-methylacetamide,
- 2-biphenyl-4-yl-N-(1-methylpyrrolidin-3-yl)-N-methylacetamide,
- 2-(3-fluorobiphenyl-4-yl)-N-(1-isopropylpyrrolidin-3-yl)-N-methylacetamide,
- 2-(2-fluorobiphenyl-4-yl)-N-(1-isopropylpyrrolidin-3-yl)-N-methylacetamide,
- 2-(2'-fluorobiphenyl-4-yl)-N-(1-isopropylpyrrolidin-3-yl)-N-methylacetamide,
- 25 2-(3'-fluorobiphenyl-4-yl)-N-(1-isopropylpyrrolidin-3-yl)-N-methylacetamide
- 2-(4'-fluorobiphenyl-4-yl)-N-(1-isopropylpyrrolidin-3-yl)-N-methylacetamide,
- N-(1-Isopropylpyrrolidin-3-yl)-N-methyl-3-(4-trifluoromethylphenyl)acrylamide

2-Biphenyl-4-yl-N-(1-isopropyl-piperidin-3-yl)-N-methyl-acetamide
2-Biphenyl-4-yl-N-(1-cyclopropylmethyl-piperidin-3-yl)-N-methyl-acetamid
N-(1-Isopropyl-piperidin-3-yl)-N-methyl-3-phenoxy-benzamide
N-(1-Cyclopropylmethyl-piperidin-3-yl)-N-methyl-3-phenoxy-benzamide
5 3-Benzo[1,3]dioxol-5-yl-N-(1-isopropyl-piperidin-3-yl)-N-methyl-acrylamide
3-Benzo[1,3]dioxol-5-yl-N-(1-cyclopropylmethyl-piperidin-3-yl)-N-methyl-acrylamide

as well as any diastereomer or enantiomer or tautomeric form thereof including mix-
tures of these or a pharmaceutically acceptable salt thereof.

10 The compounds of the present invention may be chiral, and it is intended that any
enantiomers, as separated, pure or partially purified enantiomers or racemic mixtures thereof
are included within the scope of the invention.

Furthermore, when a double bond or a fully or partially saturated ring system or
more than one center of asymmetry or a bond with restricted rotatability is present in the
15 molecule diastereomers may be formed. It is intended that any diastereomers, as separated,
pure or partially purified diastereomers or mixtures thereof are included within the scope of
the invention.

Furthermore, some of the compounds of the present invention may exist in different
tautomeric forms and it is intended that any tautomeric forms, which the compounds are able
20 to form, are included within the scope of the present invention.

The present invention also encompasses pharmaceutically acceptable salts of the
present compounds. Such salts include pharmaceutically acceptable acid addition salts,
pharmaceutically acceptable metal salts, ammonium and alkylated ammonium salts. Acid
addition salts include salts of inorganic acids as well as organic acids. Representative exam-
25 ples of suitable inorganic acids include hydrochloric, hydrobromic, hydroiodic, phosphoric,
sulfuric, nitric acids and the like. Representative examples of suitable organic acids include
formic, acetic, trichloroacetic, trifluoroacetic, propionic, benzoic, cinnamic, citric, fumaric, gly-
colic, lactic, maleic, malic, malonic, mandelic, oxalic, picric, pyruvic, salicylic, succinic,
methanesulfonic, ethanesulfonic, tartaric, ascorbic, pantoic, bismethylene salicylic, ethanedi-
30 sulfonic, gluconic, citraconic, aspartic, stearic, palmitic, EDTA, glycolic, p-aminobenzoic, glu-
tamic, benzenesulfonic, p-toluenesulfonic acids and the like. Further examples of pharma-
ceutically acceptable inorganic or organic acid addition salts include the pharmaceutically
acceptable salts listed in J. Pharm. Sci. 1977, 66, 2, which is incorporated herein by refer-
ence. Examples of metal salts include lithium, sodium, potassium, magnesium salts and the
35 like. Examples of ammonium and alkylated ammonium salts include ammonium, methylam-

monium, dimethylammonium, trimethylammonium, ethylammonium, hydroxyethylammonium, diethylammonium, butylammonium, tetramethylammonium salts and the like.

Also intended as pharmaceutically acceptable acid addition salts are the hydrates, which the present compounds are able to form.

5 The acid addition salts may be obtained as the direct products of compound synthesis. In the alternative, the free base may be dissolved in a suitable solvent containing the appropriate acid, and the salt isolated by evaporating the solvent or otherwise separating the salt and solvent.

10 The compounds of the present invention may form solvates with standard low molecular weight solvents using methods well known to the person skilled in the art. Such solvates are also contemplated as being within the scope of the present invention.

15 The invention also encompasses prodrugs of the present compounds, which on administration undergo chemical conversion by metabolic processes before becoming active pharmacological substances. In general, such prodrugs will be functional derivatives of the present compounds, which are readily convertible *in vivo* into the required compound of the formula (I). Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985.

 The invention also encompasses active metabolites of the present compounds.


20 The compounds of the present invention interact with the histamine H3 receptor and are accordingly useful for the treatment of a wide variety of conditions and disorders in which histamine H3 receptor interactions are beneficial.

25 Accordingly, in another aspect the present invention relates to a compound of the general formula (I) as well as any diastereomer or enantiomer or tautomeric form thereof including mixtures of these or a pharmaceutically acceptable salt thereof for use as a pharmaceutical composition.

30 The invention also relates to pharmaceutical compositions comprising, as an active ingredient, at least one compound of the formula (I) or any diastereomer or enantiomer or tautomeric form thereof including mixtures of these or a pharmaceutically acceptable salt thereof together with one or more pharmaceutically acceptable carriers or diluents.

Furthermore, the invention relates to the use of a compound of the general formula

(I'):



wherein

5

R¹ represents

- hydrogen,
- 10 • C₁₋₈-alkyl, C₃₋₈-alkenyl or C₃₋₈-alkynyl,
which may optionally be substituted with one or more halogen atoms,
- C₃₋₇-cycloalkyl, C₃₋₇-cycloalkenyl, C₄₋₈-bicycloalkyl, C₃₋₇-cycloalkyl-C₁₋₃-alkyl or
C₃₋₇-cycloalkenyl-C₁₋₃-alkyl,
- 15 which may optionally be substituted at any position with one or more halogen atoms,

R² represents C₁₋₆-alkyl,

X represents $-(CH_2)_m-(Z)_n-(CH_2)_o-$,

20

m and o independently are 0, 1, 2, 3 or 4,

n is 0 or 1,

25

Z is $-O-$, $-NH-$, $-N(CH_3)-$, $-C(=O)-$, $-CH(OH)-$, $-C(=N-OH)-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$, $-CH=CH-$ or $-C\equiv C-$,

Y is

(a) aryl or heteroaryl, which may optionally be substituted with one or more substituents selected from

- 5
 - halogen, nitro, cyano, hydroxy, C₁₋₆-alkanoyl, C₁₋₆-alkylthio, C₁₋₆-alkylsulfonyl, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₃₋₈-cycloalkyl, trifluoromethyl, trifluoromethoxy, -NR³R⁴ and -O(C=O)NR³R⁴, or wherein two substituents in adjacent positions form a radical -O-(CH₂)₁₋₃-O-, wherein R³ and R⁴ independently are hydrogen, C₁₋₆-alkyl, C₃₋₈-cycloalkyl, C₁₋₆-alkanoyl or aryl, or R³ and R⁴ together with the nitrogen atom to which they are attached form a 4 to 7 membered, saturated or unsaturated azetidiny, pyrrolidinyl, piperidyl or azepanyl ring,
- 10
- 15
 - aryl, aryloxy, aryl-C₁₋₆-alkyl and aryl-C₁₋₆-alkoxy, wherein the ring moieties optionally may be substituted with one or more substituents selected from
 - halogen, nitro, cyano, hydroxy, C₁₋₆-alkanoyl, C₁₋₆-alkylthio, C₁₋₆-alkylsulfonyl, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₃₋₈-cycloalkyl, trifluoromethyl, trifluoromethoxy, -NR⁵R⁶ and -O(C=O)NR⁵R⁶, or wherein two substituents in adjacent positions form a radical -O-(CH₂)₁₋₃-O-, wherein R⁵ and R⁶ independently are hydrogen, C₁₋₆-alkyl, C₃₋₈-cycloalkyl, C₁₋₆-alkanoyl or aryl, or R⁵ and R⁶ together with the nitrogen atom to which they are attached form a 4 to 7 membered, saturated or unsaturated azetidiny, pyrrolidinyl, piperidyl or azepanyl ring,
- 20
- 25 (b) C₃₋₈-cycloalkyl or C₅₋₈-cycloalkenyl, which may optionally be substituted with one or more substituents selected from
 - C₁₋₆-alkyl, C₁₋₆-alkoxy, C₁₋₆-alkylthio, cyano, trifluoromethyl, trifluoromethoxy and halogen,
- 30
 - aryl and aryloxy, wherein the ring moieties optionally may be substituted with one or more substituents selected from
 - halogen, nitro, cyano, hydroxy, C₁₋₆-alkanoyl, C₁₋₆-alkylthio, C₁₋₆-alkylsulfonyl, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₃₋₈-cycloalkyl, trifluoromethyl, trifluoromethoxy, -NR⁷R⁸ and -O(C=O)NR⁷R⁸, or wherein two substituents in adjacent positions form a radical -O-(CH₂)₁₋₃-O-,
- 35

wherein R⁷ and R⁸ independently are hydrogen, C₁₋₆-alkyl, C₃₋₆-cycloalkyl, C₁₋₆-alkanoyl or aryl, or R⁷ and R⁸ together with the nitrogen atom to which they are attached form a 4 to 7 membered, saturated or unsaturated azetidiny, pyrrolidinyl, piperidyl or azepanyl ring,

5

a well as any diastereomer or enantiomer or tautomeric form thereof including mixtures of these or a pharmaceutically acceptable salt thereof for the preparation of a pharmaceutical composition for the treatment of disorders and diseases related to the histamine H3 receptor.

10 In still another aspect, the invention relates to a method for the treatment of diseases and disorders related to the histamine H3 receptor the method comprising administering to a subject in need thereof an effective amount of a compound of the formula (I) or any diastereomer or enantiomer or tautomeric form thereof including mixtures of these or a pharmaceutically acceptable salt thereof or a pharmaceutical composition comprising the same.

15 In one aspect the invention relates to compounds with histamine H3 receptor antagonistic activity or inverse agonistic activity which may accordingly be useful in the treatment of a wide range of conditions and disorders in which histamine H3 receptor blockade is beneficial.

20 In another aspect the invention relates to compounds with histamine H3 receptor agonistic activity and which may accordingly be useful in the treatment of a wide range of conditions and disorders in which histamine H3 receptor activation is beneficial.

In one embodiment of the invention the present compounds are used for the preparation of a pharmaceutical composition for the reduction of weight.

25 In another embodiment of the invention the present compounds are used for the preparation of a pharmaceutical composition for the treatment of overweight or obesity.

In still another embodiment of the invention the present compounds are used for the preparation of a pharmaceutical composition for the suppression of appetite or satiety induction.

30 In a further embodiment of the invention the present compounds are used for the preparation of a pharmaceutical composition for the prevention and/or treatment of disorders and diseases related to overweight or obesity such as atherosclerosis, hypertension, IGT (impaired glucose tolerance), diabetes, especially type 2 diabetes (NIDDM (non-insulin dependent diabetes mellitus)), dyslipidaemia, coronary heart disease, gallbladder disease, osteoarthritis and various types of cancer such as endometrial, breast, prostate and colon cancers.

35

In yet a further embodiment of the invention the present compounds are used for the preparation of a pharmaceutical composition for the prevention and/or treatment of eating disorders such as bulimia and binge eating.

5 In a further embodiment of the invention the present compounds are used for the preparation of a pharmaceutical composition for the treatment of IGT.

In a further embodiment of the invention the present compounds are used for the preparation of a pharmaceutical composition for the treatment of type 2 diabetes. Such treatment includes inter alia treatment for the purpose of delaying or prevention of the progression from IGT to type 2 diabetes as well as delaying or prevention of the progression
10 from non-insulin requiring type 2 diabetes to insulin requiring type 2 diabetes.

The compounds of the present invention may also be used for the treatment of airway disorders such as asthma, as anti-diarrhoeals and for the modulation of gastric acid secretion.

15 Furthermore, the compounds of the present invention may be used for the treatment of diseases associated with the regulation of sleep and wakefulness and for the treatment of narcolepsy and attention deficit disorder.

Moreover, the compounds of the invention may be used as CNS stimulants or as sedatives.

20 The present compounds may also be used for the treatment of conditions associated with epilepsy. Additionally, the present compounds may be used for the treatment of motion sickness and vertigo. Furthermore, they may be useful as regulators of hypothalamo-hypophyseal secretion, antidepressants, modulators of cerebral circulation, and in the treatment of irritable bowel syndrome.

25 Further, the compounds of the present invention may be used for the treatment of dementia and Alzheimer's disease.

The compounds of the present invention may also be useful for the treatment of allergic rhinitis, ulcer or anorexia.

30 The compounds of the present invention may furthermore be useful for the treatment of migraine, see McLeod et al., *The Journal of Pharmacology and Experimental Therapeutics* **287** (1998), 43-50, and for the treatment of myocardial infarction, see Mackins et al., *Expert Opinion on Investigational Drugs* **9** (2000), 2537-2542.

In a further aspect of the invention treatment of a patient with the present compounds is combined with diet and/or exercise.

35 In a further aspect of the invention the present compounds are administered in combination with one or more further active substances in any suitable ratio(s). Such further ac-

tive agents may be selected from antiobesity agents, antidiabetics, antidyslipidemic agents, antihypertensive agents, agents for the treatment of complications resulting from or associated with diabetes and agents for the treatment of complications and disorders resulting from or associated with obesity.

5 Thus, in a further aspect of the invention the present compounds are administered in combination with one or more antiobesity agents or appetite regulating agents.

Such agents may be selected from the group consisting of CART (cocaine amphetamine regulated transcript) agonists, NPY (neuropeptide Y) antagonists, MC4 (melanocortin 4) agonists, MC3 (melanocortin 3) agonists, orexin antagonists, TNF (tumor necrosis
10 factor) agonists, CRF (corticotropin releasing factor) agonists, CRF BP (corticotropin releasing factor binding protein) antagonists, urocortin agonists, β 3 adrenergic agonists such as CL-316243, AJ-9677, GW-0604, LY362884, LY377267 or AZ-40140, MSH (melanocyte-stimulating hormone) agonists, MCH (melanocyte-concentrating hormone) antagonists, CCK (cholecystokinin) agonists, serotonin re-uptake inhibitors such as fluoxetine, seroxat or citalopram, serotonin and noradrenaline re-uptake inhibitors, mixed serotonin and noradrenergic
15 compounds, 5HT (serotonin) agonists, bombesin agonists, galanin antagonists, growth hormone, growth factors such as prolactin or placental lactogen, growth hormone releasing compounds, TRH (thyrotropin releasing hormone) agonists, UCP 2 or 3 (uncoupling protein 2 or 3) modulators, leptin agonists, DA agonists (bromocriptin, doprexin), lipase/amylase inhibitors, PPAR (peroxisome proliferator-activated receptor) modulators, RXR (retinoid X receptor) modulators, TR β agonists, AGRP (Agouti related protein) inhibitors, opioid antagonists (such as naltrexone), exendin-4, GLP-1 and ciliary neurotrophic factor.
20

In one embodiment of the invention the antiobesity agent is leptin.

In another embodiment the antiobesity agent is dexamphetamine or amphetamine.

25 In another embodiment the antiobesity agent is fenfluramine or dexfenfluramine.

In still another embodiment the antiobesity agent is sibutramine.

In a further embodiment the antiobesity agent is orlistat.

In another embodiment the antiobesity agent is mazindol or phentermine.

In still another embodiment the antiobesity agent is phendimetrazine, diethylpropion,
30 fluoxetine, bupropion, topiramate or ecopipam.

In yet another embodiment the antiobesity agent is growth hormone, a growth factor such as prolactin or placental lactogen, or a growth hormone releasing compound.

In yet a further aspect the present compounds are administered in combination with one or more antidiabetic agents.

Relevant antidiabetic agents include insulin, insulin analogues and derivatives such as those disclosed in EP 0 792 290 (Novo Nordisk A/S), eg N^{B29}-tetradecanoyl des (B30) human insulin, EP 0 214 826 and EP 0 705 275 (Novo Nordisk A/S), eg Asp^{B28} human insulin, US 5,504,188 (Eli Lilly), eg Lys^{B28} Pro^{B29} human insulin, EP 0 368 187 (Aventis), eg Lantus®, which are all incorporated herein by reference, GLP-1 derivatives such as those disclosed in WO 98/08871 (Novo Nordisk A/S), which is incorporated herein by reference, as well as orally active hypoglycaemic agents.

The orally active hypoglycaemic agents preferably comprise imidazolines, sulfonyleureas, biguanides, meglitinides, oxadiazolidinediones, thiazolidinediones, insulin sensitizers, α -glucosidase inhibitors, agents acting on the ATP-dependent potassium channel of the β -cells eg potassium channel openers such as those disclosed in WO 97/26265, WO 99/03861 and WO 00/37474 (Novo Nordisk A/S) which are incorporated herein by reference, or mitiglinide, or a potassium channel blocker, such as BTS-67582, nateglinide, glucagon antagonists such as those disclosed in WO 99/01423 and WO 00/39088 (Novo Nordisk A/S and Agouron Pharmaceuticals, Inc.), which are incorporated herein by reference, GLP-1 agonists such as those disclosed in WO 00/42026 (Novo Nordisk A/S and Agouron Pharmaceuticals, Inc.), which are incorporated herein by reference, DPP-IV (dipeptidyl peptidase-IV) inhibitors, PTPase (protein tyrosine phosphatase) inhibitors, inhibitors of hepatic enzymes involved in stimulation of gluconeogenesis and/or glycogenolysis, glucose uptake modulators, GSK-3 (glycogen synthase kinase-3) inhibitors, compounds modifying the lipid metabolism such as antilipidemic agents, compounds lowering food intake, PPAR (peroxisome proliferator-activated receptor) and RXR (retinoid X receptor) agonists, such as ALRT-268, LG-1268 or LG-1069.

In one embodiment of the invention the present compounds are administered in combination with insulin or an insulin analogue or derivative, such as N^{B29}-tetradecanoyl des (B30) human insulin, Asp^{B28} human insulin, Lys^{B28} Pro^{B29} human insulin, Lantus®, or a mixture comprising one or more of these.

In a further embodiment of the invention the present compounds are administered in combination with a sulfonyleurea eg tolbutamide, chlorpropamide, tolazamide, glibenclamide, glipizide, glimepiride, gliclazide or glyburide.

In another embodiment of the invention the present compounds are administered in combination with a biguanide eg metformin.

In yet another embodiment of the invention the present compounds are administered in combination with a meglitinide eg repaglinide or nateglinide.

In still another embodiment of the invention the present compounds are administered in combination with a thiazolidinedione insulin sensitizer eg troglitazone, ciglitazone, pioglitazone, rosiglitazone, isaglitazone, darglitazone, englitazone, CS-011/CI-1037 or T 174 or the compounds disclosed in WO 97/41097, WO 97/41119, WO 97/41120, WO 00/41121
5 and WO 98/45292 (Dr. Reddy's Research Foundation), which are incorporated herein by reference.

In still another embodiment of the invention the present compounds may be administered in combination with an insulin sensitizer eg such as GI 262570, YM-440, MCC-555, JTT-501, AR-H039242, KRP-297, GW-409544, CRE-16336, AR-H049020, LY510929,
10 MBX-102, CLX-0940, GW-501516 or the compounds disclosed in WO 99/19313, WO 00/50414, WO 00/63191, WO 00/63192, WO 00/63193 (Dr. Reddy's Research Foundation) and WO 00/23425, WO 00/23415, WO 00/23451, WO 00/23445, WO 00/23417, WO 00/23416, WO 00/63153, WO 00/63196, WO 00/63209, WO 00/63190 and WO 00/63189 (Novo Nordisk A/S), which are incorporated herein by reference.

15 In a further embodiment of the invention the present compounds are administered in combination with an α -glucosidase inhibitor eg voglibose, emiglitate, miglitol or acarbose.

In another embodiment of the invention the present compounds are administered in combination with an agent acting on the ATP-dependent potassium channel of the β -cells eg tolbutamide, glibenclamide, glipizide, gliclazide, BTS-67582 or repaglinide.

20 In yet another embodiment of the invention the present compounds may be administered in combination with nateglinide.

In still another embodiment, the present compounds are administered in combination with an antihyperlipidemic agent or antilipidemic agent, eg cholestyramine, colestipol, clofibrate, gemfibrozil, lovastatin, pravastatin, simvastatin, probucol or dextrothyroxine.

25 In still another embodiment of the invention the present compounds are administered in combination with an antilipidemic agent eg cholestyramine, colestipol, clofibrate, gemfibrozil, lovastatin, pravastatin, simvastatin, probucol or dextrothyroxine.

In another aspect of the invention, the present compounds are administered in combination with more than one of the above-mentioned compounds eg in combination with metformin and a sulfonylurea such as glyburide; a sulfonylurea and acarbose; nateglinide and metformin; acarbose and metformin; a sulfonylurea, metformin and troglitazone; insulin and a sulfonylurea; insulin and metformin; insulin, metformin and a sulfonylurea; insulin and troglitazone; insulin and lovastatin; etc.

35 Furthermore, the present compounds may be administered in combination with one or more antihypertensive agents. Examples of antihypertensive agents are β -blockers such

as alprenolol, atenolol, timolol, pindolol, propranolol and metoprolol, ACE (angiotensin converting enzyme) inhibitors such as benazepril, captopril, enalapril, fosinopril, lisinopril, quinapril and ramipril, calcium channel blockers such as nifedipine, felodipine, nicardipine, isradipine, nimodipine, diltiazem and verapamil, and α -blockers such as doxazosin, urapidil, prazosin and terazosin. Further reference can be made to Remington: The Science and Practice of Pharmacy, 19th Edition, Gennaro, Ed., Mack Publishing Co., Easton, PA, 1995.

It should be understood that any suitable combination of the compounds according to the invention with diet and/or exercise, one or more of the above-mentioned compounds and optionally one or more other active substances are considered to be within the scope of the present invention.

PHARMACEUTICAL COMPOSITIONS

The compounds of the invention may be administered alone or in combination with pharmaceutically acceptable carriers or excipients, in either single or multiple doses. The pharmaceutical compositions according to the invention may be formulated with pharmaceutically acceptable carriers or diluents as well as any other known adjuvants and excipients in accordance with conventional techniques such as those disclosed in Remington: The Science and Practice of Pharmacy, 19th Edition, Gennaro, Ed., Mack Publishing Co., Easton, PA, 1995.

The pharmaceutical compositions may be specifically formulated for administration by any suitable route such as the oral, rectal, nasal, pulmonary, topical (including buccal and sublingual), transdermal, intracisternal, intraperitoneal, vaginal and parenteral (including subcutaneous, intramuscular, intrathecal, intravenous and intradermal) route, the oral route being preferred. It will be appreciated that the preferred route will depend on the general condition and age of the subject to be treated, the nature of the condition to be treated and the active ingredient chosen.

Pharmaceutical compositions for oral administration include solid dosage forms such as capsules, tablets, dragees, pills, lozenges, powders and granules. Where appropriate, they can be prepared with coatings such as enteric coatings or they can be formulated so as to provide controlled release of the active ingredient such as sustained or prolonged release according to methods well known in the art.

Liquid dosage forms for oral administration include solutions, emulsions, suspensions, syrups and elixirs.

Pharmaceutical compositions for parenteral administration include sterile aqueous and non-aqueous injectable solutions, dispersions, suspensions or emulsions as well as ster-

ile powders to be reconstituted in sterile injectable solutions or dispersions prior to use. Depot injectable formulations are also contemplated as being within the scope of the present invention.

Other suitable administration forms include suppositories, sprays, ointments, cremes, gels, inhalants, dermal patches, implants etc.

A typical oral dosage is in the range of from about 0.001 to about 100 mg/kg body weight per day, preferably from about 0.01 to about 50 mg/kg body weight per day, and more preferred from about 0.05 to about 10 mg/kg body weight per day administered in one or more dosages such as 1 to 3 dosages. The exact dosage will depend upon the frequency and mode of administration, the sex, age, weight and general condition of the subject treated, the nature and severity of the condition treated and any concomitant diseases to be treated and other factors evident to those skilled in the art.

The formulations may conveniently be presented in unit dosage form by methods known to those skilled in the art. A typical unit dosage form for oral administration one or more times per day such as 1 to 3 times per day may contain of from 0.05 to about 1000 mg, preferably from about 0.1 to about 500 mg, and more preferred from about 0.5 mg to about 200 mg.

For parenteral routes, such as intravenous, intrathecal, intramuscular and similar administration, typically doses are in the order of about half the dose employed for oral administration.

The compounds of this invention are generally utilized as the free substance or as a pharmaceutically acceptable salt thereof. One example is an acid addition salt of a compound having the utility of a free base. When a compound of the formula (I) contains a free base such salts are prepared in a conventional manner by treating a solution or suspension of a free base of the formula (I) with a chemical equivalent of a pharmaceutically acceptable acid, for example, inorganic and organic acids. Representative examples are mentioned above. Physiologically acceptable salts of a compound with a hydroxy group include the anion of said compound in combination with a suitable cation such as sodium or ammonium ion.

For parenteral administration, solutions of the novel compounds of the formula (I) in sterile aqueous solution, aqueous propylene glycol or sesame or peanut oil may be employed. Such aqueous solutions should be suitable buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. The aqueous solutions are particularly suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administra-

tion. The sterile aqueous media employed are all readily available by standard techniques known to those skilled in the art.

Suitable pharmaceutical carriers include inert solid diluents or fillers, sterile aqueous solution and various organic solvents. Examples of solid carriers are lactose, terra alba, sucrose, cyclodextrin, talc, gelatine, agar, pectin, acacia, magnesium stearate, stearic acid or lower alkyl ethers of cellulose. Examples of liquid carriers are syrup, peanut oil, olive oil, phospholipids, fatty acids, fatty acid amines, polyoxyethylene or water. Similarly, the carrier or diluent may include any sustained release material known in the art, such as glyceryl monostearate or glyceryl distearate, alone or mixed with a wax. The pharmaceutical compositions formed by combining the novel compounds of the formula (I) and the pharmaceutically acceptable carriers are then readily administered in a variety of dosage forms suitable for the disclosed routes of administration. The formulations may conveniently be presented in unit dosage form by methods known in the art of pharmacy.

Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules or tablets, each containing a predetermined amount of the active ingredient, and which may include a suitable excipient. These formulations may be in the form of powder or granules, as a solution or suspension in an aqueous or non-aqueous liquid, or as an oil-in-water or water-in-oil liquid emulsion.

If a solid carrier is used for oral administration, the preparation may be tableted, placed in a hard gelatine capsule in powder or pellet form or it can be in the form of a troche or lozenge. The amount of solid carrier will vary widely but will usually be from about 25 mg to about 1 g. If a liquid carrier is used, the preparation may be in the form of a syrup, emulsion, soft gelatine capsule or sterile injectable liquid such as an aqueous or non-aqueous liquid suspension or solution.

A typical tablet, which may be prepared by conventional tableting techniques, may contain:

Core:		
	Active compound (as free compound or salt thereof)	5.0 mg
	Lactosum Ph. Eur.	67.8 mg
30	Cellulose, microcryst. (Avicel)	31.4 mg
	Amberlite® IRP88*	1.0 mg
	Magnesii stearas Ph. Eur.	q.s.

Coating:

Hydroxypropyl methylcellulose	approx.	9 mg
Mywacett 9-40 T**	approx.	0.9 mg

5 * Polacrillin potassium NF, tablet disintegrant, Rohm and Haas.

 ** Acylated monoglyceride used as plasticizer for film coating.

 If desired, the pharmaceutical composition of the invention may comprise the compound of the formula (I) in combination with further pharmacologically active substances
 10 such as those described in the foregoing.

EXAMPLES


 In the examples the following terms are intended to have the following, general meanings:

- DCM: dichloromethane
 15 DIPEA: diisopropylethylamine
 DMF: *N,N*-dimethylformamide
 DMSO: dimethyl sulphoxide
 THF: tetrahydrofuran

 NMR spectra were recorded on Bruker 300 MHz and 400 MHz instruments. HPLC-
 20 MS was performed on a Perkin Elmer instrument (API 100). The column used was X-Terra C-18, 5 μ m, 50 X 3 mm, and elution was done at 1.5 ml/min at room temperature with a gradient of 5% to 90% acetonitrile in water with 0.01% trifluoroacetic acid within 7.5 min.

General Procedure (A)

 The compounds of formula (I) can be prepared according to the following general
 25 procedure (A):




 To a mixture of the acid (150 mmol), DMF (200 ml), and *N*-hydroxybenzotriazole (40.6 g, 301
 30 mmol) is added a solution of *N*-ethyl-*N*-(3-dimethylaminopropyl)carbodiimide hydrochloride (28.8 g, 150 mmol) in DMF (100 ml). The mixture was stirred at room temperature for 1.5

hour, and a solution of the diamine (150 mmol) in DCM (100 ml) is added. The mixture is stirred at room temperature for 4 hours, concentrated under reduced pressure, and the residue is distributed between ethyl acetate (1.0 l) and a saturated, aqueous NaHCO₃ solution (1.0 l). The phases are separated, the organic layer is dried (MgSO₄), and concentrated, and the residue is re-dissolved in 1 M aqueous hydrochloric acid (150 ml). The solution is concentrated, and the residue is dried by co-evaporation with ethanol. Re-crystallization of the residue from ethanol yielded the compound of formula (Ia).

Example 1 (general procedure (A))


4-(3,4-Dimethoxyphenyl)-*N*-[1-(1-ethylpropyl)pyrrolidin-3-yl]-*N*-methyl-4-oxobutamide hydrochloride




¹H NMR (DMSO-*d*₆) δ 0.92 (m, 6H), 1.70 (m, 4H), 1.95-2.30 (m, 2H), 2.52-3.65 (m, 12H), 3.80 (s, 3H), 3.84 (s, 3H), 4.90-5.20 (m, 1H), 7.07 (d, *J* = 8 Hz, 1H), 7.45 (br s, 1H), 7.68 (br d, *J* = 8 Hz, 1H), 10.70 (m, 1H); HPLC-MS: *m/z* 391 (MH⁺); R_f: 2.40 min.

Example 2 (general procedure (A))


2-(4-Dimethylaminophenyl)-*N*-[1-(1-ethylpropyl)pyrrolidin-3-yl]-*N*-methylacetamide hydrochloride




¹H NMR (DMSO-*d*₆) δ 0.90 (m, 6H), 1.70 (m, 4H), 1.95-2.18 (m, 2H), 2.75-3.90 (m, 16H), 4.90-5.20 (m, 1H), 7.27-7.60 (m, 4H), 10.40-11.20 (m, 1H); HPLC-MS: *m/z* 332 (MH⁺); R_f: 1.73 min.

Exempl 3 (gen ral procedure (A))2-Biphenyl-4-yl-*N*-[1-(1-ethylpropyl)pyrrolidin-3-yl]-*N*-methylacetamide hydrochloride

¹H NMR (DMSO-*d*₆) δ 0.90 (m, 6H), 1.70 (m, 4H), 1.96-2.20 (m, 2H), 2.75-3.90 (m, 10H),
5 4.90-5.15 (m, 1H), 7.29-7.40 (m, 3H), 7.46 (t, *J* = 8 Hz, 2H), 7.59-7.68 (m, 4H), 10.10-10.90
(m, 1H); HPLC-MS: *m/z* 365 (MH⁺); R_f: 3.77 min.

Example 4 (general procedure (A))4-(4-Chlorophenyl)-*N*-[1-(1-ethylpropyl)pyrrolidin-3-yl]-*N*-methyl-4-oxobutyr-
amide hydrochloride


¹H NMR (DMSO-*d*₆) δ 0.90 (m, 6H), 1.70 (m, 4H), 1.95-2.20 (m, 2H), 2.67-3.63 (m, 12H),
4.85-5.15 (m, 1H), 7.59 (d, *J* = 8 Hz, 2H), 7.99 (d, *J* = 8 Hz, 2H), 10.3-11.1 (m, 1H); HPLC-
MS: *m/z* 367 (MH⁺); R_f: 3.00 min.

Example 5 (general procedure (A))15 *N*-[1-(1-Ethylpropyl)pyrrolidin-3-yl]-*N*-methyl-3-(4-trifluoromethylphenyl)propionamide hydro-
chloride

^1H NMR (DMSO- d_6) δ 0.90 (m, 6H), 1.70 (m, 4H), 1.90-2.20 (m, 2H), 2.67-3.63 (m, 12H), 4.78-5.22 (m, 1H), 7.49 (m, 2H), 7.63 (d, J = 8 Hz, 2H), 10.4-11.1 (m, 1H); HPLC-MS: m/z 372 (MH^+); Rf: 3.50 min.

Example 6 (general procedure (A))


5 *N*-[1-(1-Ethylpropyl)pyrrolidin-3-yl]-*N*-methyl-3-phenoxybenzamide hydrochloride



^1H NMR (DMSO- d_6) δ 0.90 (m, 6H), 1.70 (m, 4H), 2.00-2.30 (m, 2H), 2.85-3.70 (m, 8H), 4.45-5.20 (m, 1H), 6.90-7.21 (m, 6H), 7.40-7.50 (m, 3H), 10.2-11.0 (m, 1H); HPLC-MS: m/z 368 (MH^+); Rf: 3.33 min.

10 **Example 7** (general procedure (A))


4-(3,4-Dimethoxyphenyl)-*N*-[1-(1-ethylpropyl)pyrrolidin-3-yl]-4-oxobutamide hydrochloride



15 ^1H NMR (DMSO- d_6) δ 0.93 (br t, J = 7 Hz, 6H), 1.69 (m, 4H), 1.90 (m, 1H), 2.10-2.35 (m, 1H), 2.47 (t, J = 7 Hz, 2H), 2.92 (m, 0.5H), 3.12 (m, 1H), 3.23 (t, J = 7 Hz, 2H), 3.25-3.70 (m, 3.5H), 3.81 (s, 3H), 3.84 (s, 3H), 4.35 (m, 1H), 7.07 (d, J = 8 Hz, 1H), 7.44 (m, 1H), 7.68 (dd, J = 1 Hz, 8 Hz, 1H), 8.41 (d, J = 7 Hz, 0.5 H), 8.52 (d, J = 7 Hz, 0.5 H), 10.35 (m, 1H); HPLC-MS: m/z 377 (MH^+); Rf: 1.79 min.

Example 8 (general procedure (A))


N-[1-(1-Ethylpropyl)pyrrolidin-3-yl]-2-(3-fluoro-4-hydroxyphenyl)-*N*-methylacetamide hydrochloride



- 5 ^1H NMR (DMSO- d_6) δ 0.90 (m, 6H), 1.70 (m, 4H), 1.90-2.20 (m, 2H), 2.70-3.25 (m, 10H), 4.85-5.15 (m, 1H), 6.78-7.10 (m, 3H), 9.70 (br s, 1H), 10.25-11.05 (m, 1H); HPLC-MS: m/z 323 (MH^+); Rf: 1.63 min.

Example 9 (general procedure (A))

2-Biphenyl-4-yl-*N*-[1-(1-ethylpropyl)pyrrolidin-3-yl]-*N*-methylacetamide hydrochloride




10

- ^1H NMR (DMSO- d_6) δ 0.38 (m, 2H), 0.59 (m, 2H), 1.07 (m, 1H), 2.00-2.25 (m, 2H), 2.70-4.00 (m, 11H), 4.88-5.09 (m, 1H), 7.29-7.38 (m, 3H), 7.46 (t, $J = 8$ Hz, 2H), 7.59-7.66 (m, 4H), 10.55-11.30 (m, 1H); HPLC-MS: m/z 349 (MH^+); Rf: 2.94 min.

Example 10 (general procedure (A))


- 15 2-Biphenyl-4-yl-*N*-(1-cyclopentylpyrrolidin-3-yl)-*N*-methylacetamide hydrochloride



^1H NMR ($\text{DMSO}-d_6$) δ 1.53 (m, 2H), 1.72 (m, 4H), 1.85-2.19 (m, 4H), 2.70-4.00 (m, 10H), 4.85-5.29 (m, 1H), 7.29-7.38 (m, 3H), 7.46 (t, $J = 8$ Hz, 2H), 7.59-7.66 (m, 4H), 10.90-11.60 (m, 1H); HPLC-MS: m/z 363 (MH^+); Rf: 2.86 min.

Example 11 (general procedure (A))


- 5 3-Benzo[1,3]dioxol-5-yl-*N*-[1-(1-ethylpropyl)pyrrolidin-3-yl]-*N*-methylacrylamide hydrochloride



^1H NMR ($\text{DMSO}-d_6$) δ 0.90 (m, 6H), 1.73 (m, 4H), 2.05-2.25 (m, 2H), 2.80-4.05 (m, 8H), 5.05-5.22 (m, 1H), 6.07 (s, 2H), 6.91-7.18 (m, 3H), 7.40-7.47 (m, 2H), 10.40-11.20 (m, 1H); HPLC-MS: m/z 345 (MH^+); Rf: 2.36 min.

- 10 **Example 12** (general procedure (A))

3-(3-Chlorophenyl)-*N*-[1-(1-ethylpropyl)pyrrolidin-3-yl]-*N*-methylacrylamide hydrochloride




^1H NMR ($\text{DMSO}-d_6$) δ 0.90 (m, 6H), 1.73 (m, 4H), 2.05-2.30 (m, 2H), 2.85-3.70 (m, 8H), 5.05-5.22 (m, 1H), 7.21-7.52 (m, 4H), 7.68 (br s, 1H), 7.89 (br s, 1H), 10.20-11.05 (m, 1H); HPLC-MS: m/z 335 (MH^+); Rf: 3.07 min.

15

Example 13 (general procedure (A))


2-Biphenyl-4-yl-*N*-(1-isopropylpyrrolidin-3-yl)-*N*-methylacetamide hydrochloride



^1H NMR ($\text{DMSO}-d_6$) δ 1.28 (d, $J = 7$ Hz, 6H), 1.95-2.20 (m, 2H), 2.76-3.87 (m, 10H), 4.92-5.25 (m, 1H), 7.29-7.38 (m, 3H), 7.46 (t, $J = 8$ Hz, 2H), 7.59-7.66 (m, 4H), 10.80-11.50 (m, 1H); HPLC-MS: m/z 337 (MH^+); Rf: 3.07 min.

Example 14 (general procedure (A))


- 5 2-Biphenyl-4-yl-*N*-(1-cyclobutylpyrrolidin-3-yl)-*N*-methylacetamide hydrochloride



^1H NMR ($\text{DMSO}-d_6$) δ 1.64-1.86 (m, 2H), 1.95-2.40 (m, 6H), 2.70-3.85 (m, 10H), 4.80-5.25 (m, 1H), 7.29-7.38 (m, 3H), 7.46 (t, $J = 8$ Hz, 2H), 7.59-7.66 (m, 4H), 11.20-11.80 (m, 1H); HPLC-MS: m/z 349 (MH^+); Rf: 3.07 min.

- 10 **Example 15** (general procedure (A))


2-Biphenyl-4-yl-*N*-(1-methylpyrrolidin-3-yl)-*N*-methylacetamide hydrochloride



- 15 ^1H NMR ($\text{DMSO}-d_6$) δ 1.93-2.30 (m, 2H), 2.73-2.87 (m, 3H), 2.90-3.68 (m, 7H), 3.70-3.90 (m, 2H), 4.85-5.15 (m, 1H), 7.32-7.40 (m, 3H), 7.44 (t, $J = 8$ Hz, 2H), 7.59-7.66 (m, 4H), 10.80-11.60 (m, 1H); HPLC-MS: m/z 309 (MH^+); Rf: 3.27 min.

Example 16 (general procedure (A))

2-(2-Fluorobiphenyl-4-yl)-*N*-(1-isopropylpyrrolidin-3-yl)-*N*-methylacetamide hydrochloride




^1H NMR (DMSO- d_6) δ 1.28 (d, $J = 7$ Hz, 6H), 2.09 (m, 2H), 2.82 (m, 1H), 3.04 (m, 4H), 3.20-3.63 (m, 3H), 3.78-3.88 (m, 2H), 4.93 (m, 0.5 H), 5.18 (m, 0.5H), 7.25-7.55 (m, 8H), 10.70-11.45 (m, 1H); HPLC-MS: m/z 355 (MH^+); Rf: 3.60 min.

5

Example 17 (general procedure (A))

N-(1-isopropylpyrrolidin-3-yl)-N-methyl-3-(4-trifluoromethylphenyl)acrylamide hydrochloride




10

^1H NMR (DMSO- d_6) δ 1.31 (d, $J = 7$ Hz, 6H), 2.13 (m, 2H), 2.88-3.70 (m, 8H), 5.00-5.40 (m, 1H), 7.28-7.61 (m, 2H), 7.78 (m, 2H), 7.97 (m, 2H), 10.95-11.75 (m, 1H); HPLC-MS: m/z 341 (MH^+); Rf: 3.47 min.

15

Example 18 (general procedure (A))

2-Biphenyl-4-yl-N-(1-isopropyl-piperidin-3-yl)-N-methyl-acetamide hydrochloride



20

^1H NMR (DMSO- d_6) δ 1.26 (d, $J = 7$ Hz, 3H), 1.30 (d, $J = 7$ Hz, 3H), 1.38 (m, 1H), 2.7-3.55 (m, 11H), 3.65-4.05 (m, 2H), 4.75 (m, 1H), 7.34 (dd, 2H), 7.4-7.54 (m, 3H), 7.55-7.69 (m, 4H), 10.03 (s, 0.5H), 11.09 (s, 0.5H);

C₂₃ H₃₀ N₂ O, HCl + 1.25 mol H₂O


Calc.: C 67.15 H 8.18 N 6.54

Found.: C 67.46 H 8.25 N 6.84.

5

Example 19 (general procedure (A))

2-Biphenyl-4-yl-N-(1-cyclopropylmethyl-piperidin-3-yl)-N-methyl-acetamide
hydrochloride




10

¹H NMR (DMSO-*d*₆) δ 0.38 (m, 2H), 0.63 (m, 2H), 1.14 (m, 1H), 1.35-2.02 (m, 4H), 2.65-3.55 (m, 9H), 3.71-4.02 (m, 2H), 4.70 (m, 1H), 7.33 (dd, 2H), 7.42-7.52 (m, 3H), 7.57-7.68 (m, 4H), 10.38 (broad s, 0.5H), 11.30 (broad s, 0.5H); HPLC-MS: *m/z* 363 (MH⁺); Rf: 3.70 min.

15

Example 20 (general procedure (A))

N-(1-Isopropyl-piperidin-3-yl)-N-methyl-3-phenoxy-benzamide hydrochloride




20

¹H NMR (CDCl₃) of the free base δ 1.00 (m, 6H), 1.3-1.9 (m, 4H), 1.97 (t, 1H), 2.22 (t, 1H), 2.6-3.0 (m, 6H), 3.60-3.83 (m, 1H), 6.9-7.16 (m, 6H), 7.30-7.40 (m, 3H); HPLC-MS: *m/z* 353 (MH⁺); Rf: 2.78 min.

Example 21 (general procedure (A))


N-(1-Cyclopropylmethyl-piperidin-3-yl)-N-methyl-3-phenoxy-benzamide hydrochloride



^1H NMR (CDCl_3) δ 0.45 (m, 2H), 0.81 (m, 2H), 1.30 (m, 1H), 2.02 (m, 2H), 2.2-2.55 (m, 3H), 2.71 (m, 1H), 2.93 (dd, 2H), 3.06 (s, 3H), 3.63 (m, 2H), 3.88 (m, 0.5 H), 4.15 (m, 0.5 H), 6.95-7.10 (m, 5H), 7.15 (t, 1H), 7.33-7.42 (m, 3H), 12.10 (broad s, 1H); HPLC-MS: m/z 365 (MH^+); Rf: 2.80 min.

Example 22 (general procedure (A))


3-Benzo[1,3]dioxol-5-yl-N-(1-isopropyl-piperidin-3-yl)-N-methyl-acrylamide hydrochloride



^1H NMR (CDCl_3) δ 1.45 (m, 6H), 1.8-2.15 (m, 4H), 2.35-2.85 (m, 3H), 2.95 (m, 1H), 3.20-3.55 (m, 4H), 3.96 (m, 0.5H), 4.15 (m, 0.5H), 5.38 (m, 0.5H), 6.00 (broad s, 2H), 6.60 (m, 0.5H), 6.81 (d, $J = 8$ Hz, 1H), 7.02 (m, 1H), 7.27-7.75 (m, 2H), 11.85 (broad s, 0.5H), 12.75 (broad s, 0.5H); HPLC-MS: m/z 331 (MH^+); Rf: 2.11 min.

Example 23 (general procedure (A))

3-Benzo[1,3]dioxol-5-yl-N-(1-cyclopropylmethyl-piperidin-3-yl)-N-methyl-acrylamide hydrochloride



^1H NMR (CDCl_3) δ 0.45 (m, 2H), 0.80 (d, 2H), 1.31 (m, 1H), 1.70-2.15 (m, 4H), 2.27-3.36 (m, 8H), 3.38-4.15 (m, 3H), 5.23 (m, 0.5H), 6.00 (m, 2H), 6.60 (m, 0.5H), 6.81 (d, $J = 8\text{ Hz}$, 1H), 6.94-7.09 (m, 1H), 7.27-7.75 (m, 2H), 12.13 (broad s, 0.5H), 12.97 (broad s, 0.5H);

HPLC-MS: m/z 343 (MH^+); Rf: 2.22 min.

PHARMACOLOGICAL METHODS

The ability of the compounds to interact with the histamine H3 receptor can be determined by the following *in vitro* binding assays.

Binding assay I

The H3-receptor agonist ligand R- α -methyl[^3H]histamine (RAMHA) is incubated with isolated rat cortex cell-membranes at 25 °C for 1 hour, followed by a filtration of the incubate through Whatman GF/B filters. Radioactivity retained on the filters is measured using a beta counter.

Male Wistar rats (150-200 g) are decapitated and cerebral cortex is quickly dissected out and frozen immediately on dry ice. Tissue is kept at -80 °C until membrane preparation. During the membrane preparation the tissue is kept on ice all the time. Rat cerebral cortex is homogenized in 10 volumes (w/w) ice-cold Hepes buffer (20 mM Hepes, 5 mM MgCl_2 pH 7.1 (KOH) + 1 mg/ml bacitracin) using an Ultra-Turrax homogenizer for 30 seconds. The homogenate is centrifuged at 140 g in 10 min. The supernatant is transferred to a new test tube and centrifuged for 30 min at 30 000 g. Pellet is resuspended in 5-10 ml Hepes buffer, homogenized and centrifuged for 10 min at 30 000 g. This short centrifugation step is repeated twice. After the last centrifugation the pellet is resuspended in 2-4 ml Hepes buffer and the protein concentration is determined. The membranes are diluted to a protein concentration of 5 mg/ml using Hepes buffer, aliquoted and stored at -80 °C until use.

50 µl test-compound, 100 µl membrane (200 µg/ml), 300 µl Hepes buffer and 50 µl R-α-methyl[³H]histamine (1 nM) are mixed in a test tube. The compounds to be tested are dissolved in DMSO and further diluted in H₂O to the desired concentrations. Radioligand and membranes are diluted in Hepes buffer + 1 mg/ml bacitracin. The mixture is incubated for 60 min at 25 °C. Incubation is terminated by adding 5 ml ice-cold 0.9% NaCl, followed by rapid filtration through Whatman GF/B filters pre-treated for 1 hour with 0.5% polyethyleneimine. The filters are washed with 2 x 5 ml ice-cold NaCl. To each filter a 3 ml scintillation cocktail is added and the radioactivity retained is measured with a Packard Tri-Carb beta counter.

IC₅₀ values are calculated by non-linear regression analysis of binding curves (6 points minimum) using the windows program GraphPad Prism, GraphPad software, USA.

Binding assay II

The human H3 receptor is cloned by PCR and subcloned into the pcDNA3 expression vector. Cells stably expressing the H3 receptor are generated by transfecting the H3-expression vectors into HEK 293 cells and using G418 to select for H3 clones. The human H3-HEK 293 clones are cultured in DMEM (GIBCO-BRL) with glutamax, 10% foetal calf serum, 1% penicillin/streptavidin and 1 mg/ml G 418 at 37 °C and 5% CO₂. Before harvesting, the confluent cells are rinsed with PBS and incubated with Versene (proteinase, GIBCO-BRL) for approximately 5 min. The cells are flushed with PBS and DMEM and the cell suspension collected in a tube and centrifuged for 5-10 min at 420 g in a Heraeus Sepatech Megafuge 1.0. The pellet is resuspended in 10-20 vol. Hepes buffer (20 mM Hepes, 5 mM MgCl₂, pH 7.1 (KOH)) and homogenized for 10-20 seconds using an Ultra-Turrax homogenizer. The homogenate is centrifuged for 30 min at 30 000 g. The pellet is resuspended in 5-10 ml Hepes buffer, homogenized 5-10 seconds with the Ultra-Turrax and centrifuged for 10 min at 30 000 g. Following this centrifugation step, the membrane pellet is resuspended in 2-4 ml Hepes buffer, homogenized with a syringe or Teflon homogenizer, and the protein concentration determined. The membranes are diluted to a protein concentration of 1-5 mg/ml in Hepes buffer, aliquoted and kept at -80 °C until use.

Aliquots of the membrane suspension are incubated for 60 min at 25 °C with 30 pM [¹²⁵I]-iodoproxifan, a known compound with high affinity for the H3 receptor, and the test compound at various concentrations. The incubation is stopped by dilution with ice-cold medium, followed by rapid filtration through Whatman GF/B filters pretreated for 1 hour with 0.5% polyethyleneimine. The radioactivity retained on the filters is counted using a Cobra II auto gamma counter. The radioactivity of the filters is indirectly proportional to the binding affinity of the tested compound. The results are analysed by nonlinear regression analysis.

When tested, the present compounds of the formula (I) generally show a high binding affinity to the histamine H3 receptor.

Functional assay I

5 The ability of the compounds to interact with the histamine H3 receptor as agonists, inverse agonists and/or antagonists, is determined by an *in vitro* functional assay utilizing membranes from HEK 293 cell expressing the human H3 receptors.

The H3 receptor is cloned by PCR and subcloned into the pcDNA3 expression vector. Cells stably expressing the H3 receptor are generated by transfecting the H3-expression vectors into HEK 293 cells and using G418 to select for H3 clones. The human H3-HEK 293
10 clones are cultured in DMEM with glutamax, 10% foetal calf serum, 1% penicillin/streptavidin and 1 mg/ml G418 at 37 °C and 5% CO₂.

The H3 receptor expressing cells are washed once with phosphate buffered saline (PBS) and harvested using versene (GIBCO-BRL). PBS is added and the cells are centrifuged for 5 min at 364 g. The cell pellet is resuspended in stimulation buffer to a concentration of 1 x 10⁶ cells/ml. cAMP accumulation is measured using the Flash Plate® cAMP assay (NEN™ Life Science Products). The assay is generally performed as described by the manufacturer. Briefly, 50 µl cell suspension is added to each well of the Flashplate which also contained 25 µl 40 µM isoprenaline, to stimulate cAMP generation, and 25 µl of test compound (either agonists or inverse agonists alone, or agonist and antagonist in combination). The
15 assay can be run in "agonist-mode" which means that the test compound is added, in increasing concentration, on its own, to the cells, and cAMP is measured. If cAMP goes up, it is an inverse agonist; if cAMP does not change, it is a neutral antagonist, and if cAMP goes down, it is an agonist. The assay can also be run in the "antagonist-mode" which means that a test compound is added, in increasing concentrations, together with increasing concentrations of a known H3 agonist (eg RAMHA). If the compound is an antagonist, increasing concentrations of it cause a right-ward shift in the H3 agonist's dose-response curves. The final
25 volume in each well is 100 µl. Test compounds are dissolved in DMSO and diluted in H₂O. The mixture is shaken for 5 min, and allowed to stand for 25 min at room temperature. The reaction is stopped with 100 µl "Detection Mix" per well. The plates are then sealed with plastic,
30 shaken for 30 min, allowed to stand overnight, and finally the radioactivity is counted in the Cobra II auto gamma topcounter. EC₅₀ values are calculated by non-linear regression analysis of dose response curves (6 points minimum) using GraphPad Prism. Kb values are calculated by Schild plot analysis.

Functional assay II

The ability of the compounds to bind and interact with the human, monkey or rat H3 receptor as agonists, inverse agonists and/or antagonists, is determined by a functional assay, named [³⁵S] GTP γ S assay.

5 The human H3 receptor has the following sequence:

Met-Glu-Arg-Ala-Pro-Pro-Asp-Gly-Pro-Leu-Asn-Ala-Ser-Gly-Ala-Leu-Ala-
Gly-Glu-Ala-Ala-Ala-Ala-Gly-Gly-Ala-Arg-Gly-Phe-Ser-Ala-Ala-Trp-Thr-
Ala-Val-Leu-Ala-Ala-Leu-Met-Ala-Leu-Leu-Ile-Val-Ala-Thr-Val-Leu-Gly-
Asn-Ala-Leu-Val-Met-Leu-Ala-Phe-Val-Ala-Asp-Ser-Ser-Leu-Arg-Thr-Gln-
10 Asn-Asn-Phe-Phe-Leu-Leu-Asn-Leu-Ala-Ile-Ser-Asp-Phe-Leu-Val-Gly-Ala-
Phe-Cys-Ile-Pro-Leu-Tyr-Val-Pro-Tyr-Val-Leu-Thr-Gly-Arg-Trp-Thr-Phe-
Gly-Arg-Gly-Leu-Cys-Lys-Leu-Trp-Leu-Val-Val-Asp-Tyr-Leu-Leu-Cys-Thr-
Ser-Ser-Ala-Phe-Asn-Ile-Val-Leu-Ile-Ser-Tyr-Asp-Arg-Phe-Leu-Ser-Val-
Thr-Arg-Ala-Val-Ser-Tyr-Arg-Ala-Gln-Gln-Gly-Asp-Thr-Arg-Arg-Ala-Val-
15 Arg-Lys-Met-Leu-Leu-Val-Trp-Val-Leu-Ala-Phe-Leu-Leu-Tyr-Gly-Pro-Ala-
Ile-Leu-Ser-Trp-Glu-Tyr-Leu-Ser-Gly-Gly-Ser-Ser-Ile-Pro-Glu-Gly-His-
Cys-Tyr-Ala-Glu-Phe-Phe-Tyr-Asn-Trp-Tyr-Phe-Leu-Ile-Thr-Ala-Ser-Thr-
Leu-Glu-Phe-Phe-Thr-Pro-Phe-Leu-Ser-Val-Thr-Phe-Phe-Asn-Leu-Ser-Ile-
Tyr-Leu-Asn-Ile-Gln-Arg-Arg-Thr-Arg-Leu-Arg-Leu-Asp-Gly-Ala-Arg-Glu-
20 Ala-Ala-Gly-Pro-Glu-Pro-Pro-Pro-Glu-Ala-Gln-Pro-Ser-Pro-Pro-Pro-Pro-
Pro-Gly-Cys-Trp-Gly-Cys-Trp-Gln-Lys-Gly-His-Gly-Glu-Ala-Met-Pro-Leu-
His-Arg-Tyr-Gly-Val-Gly-Glu-Ala-Ala-Val-Gly-Ala-Glu-Ala-Gly-Glu-Ala-
Thr-Leu-Gly-Gly-Gly-Gly-Gly-Gly-Gly-Ser-Val-Ala-Ser-Pro-Thr-Ser-Ser-
Ser-Gly-Ser-Ser-Ser-Arg-Gly-Thr-Glu-Arg-Pro-Arg-Ser-Leu-Lys-Arg-Gly-
25 Ser-Lys-Pro-Ser-Ala-Ser-Ser-Ala-Ser-Leu-Glu-Lys-Arg-Met-Lys-Met-Val-
Ser-Gln-Ser-Phe-Thr-Gln-Arg-Phe-Arg-Leu-Ser-Arg-Asp-Arg-Lys-Val-Ala-
Lys-Ser-Leu-Ala-Val-Ile-Val-Ser-Ile-Phe-Gly-Leu-Cys-Trp-Ala-Pro-Tyr-
Thr-Leu-Leu-Met-Ile-Ile-Arg-Ala-Ala-Cys-His-Gly-His-Cys-Val-Pro-Asp-
Tyr-Trp-Tyr-Glu-Thr-Ser-Phe-Trp-Leu-Leu-Trp-Ala-Asn-Ser-Ala-Val-Asn-
30 Pro-Val-Leu-Tyr-Pro-Leu-Cys-His-His-Ser-Phe-Arg-Arg-Ala-Phe-Thr-Lys-
Leu-Leu-Cys-Pro-Gln-Lys-Leu-Lys-Ile-Gln-Pro-His-Ser-Ser-Leu-Glu-His-
Cys-Trp-Lys

The monkey H3 receptor has the following sequence:

Met-Glu-Arg-Ala-Pro-Pro-Asp-Gly-Pro-Leu-Asn-Ala-Ser-Gly-Ala-Leu-Ala-
35 Gly-Glu-Ala-Ala-Ala-Ala-Gly-Gly-Ala-Arg-Gly-Phe-Ser-Ala-Ala-Trp-Thr-

Ala-Val-Leu-Ala-Ala-Leu-Met-Ala-Leu-Leu-Ile-Val-Ala-Thr-Val-Leu-Gly-
 Asn-Ala-Leu-Val-Met-Leu-Ala-Phe-Val-Ala-Asp-Ser-Ser-Leu-Arg-Thr-Gln-
 Asn-Asn-Phe-Phe-Leu-Leu-Asn-Leu-Ala-Ile-Ser-Asp-Phe-Leu-Val-Gly-Ala-
 Phe-Cys-Ile-Pro-Leu-Tyr-Val-Pro-Tyr-Val-Leu-Thr-Gly-Arg-Trp-Thr-Phe-
 5 Gly-Arg-Gly-Leu-Cys-Lys-Leu-Trp-Leu-Val-Val-Asp-Tyr-Leu-Leu-Cys-Thr-
 Ser-Ser-Ala-Phe-Asn-Ile-Val-Leu-Ile-Ser-Tyr-Asp-Arg-Phe-Leu-Ser-Val-
 Thr-Arg-Ala-Val-Ser-Tyr-Arg-Ala-Gln-Gln-Gly-Asn-Thr-Arg-Arg-Ala-Val-
 Arg-Lys-Met-Leu-Leu-Val-Trp-Val-Leu-Ala-Phe-Leu-Leu-Tyr-Gly-Pro-Ala-
 Ile-Leu-Ser-Trp-Glu-Tyr-Leu-Ser-Gly-Gly-Ser-Ser-Ile-Pro-Glu-Gly-His-
 10 Cys-Tyr-Ala-Glu-Phe-Phe-Tyr-Asn-Trp-Tyr-Phe-Leu-Ile-Thr-Ala-Ser-Thr-
 Leu-Glu-Phe-Phe-Thr-Pro-Phe-Leu-Ser-Val-Thr-Phe-Phe-Asn-Leu-Ser-Ile-
 Tyr-Leu-Asn-Ile-Gln-Arg-Arg-Thr-Arg-Leu-Arg-Leu-Asp-Gly-Ala-Arg-Glu-
 Ala-Gly-Gly-Pro-Glu-Pro-Pro-Pro-Glu-Ala-Gln-Pro-Ser-Pro-Pro-Pro-Pro-
 Pro-Gly-Cys-Trp-Gly-Cys-Trp-Gln-Lys-Gly-His-Gly-Glu-Ala-Met-Pro-Leu-
 15 His-Arg-Tyr-Gly-Val-Gly-Glu-Ala-Ala-Ala-Gly-Ala-Glu-Ala-Gly-Glu-Thr-
 Ala-Leu-Gly-Gly-Gly-Gly-Gly-Gly-Gly-Ser-Ala-Ala-Ser-Pro-Thr-Ser-Ser-
 Ser-Gly-Ser-Ser-Ser-Arg-Gly-Thr-Glu-Arg-Pro-Arg-Ser-Leu-Lys-Arg-Gly-
 Ser-Lys-Pro-Ser-Ala-Ser-Ser-Ala-Ser-Leu-Glu-Lys-Arg-Met-Lys-Met-Val-
 Ser-Gln-Ser-Phe-Thr-Gln-Arg-Phe-Arg-Leu-Ser-Arg-Asp-Arg-Lys-Val-Ala-
 20 Lys-Ser-Leu-Ala-Val-Ile-Val-Ser-Ile-Phe-Gly-Leu-Cys-Trp-Ala-Pro-Tyr-
 Thr-Leu-Leu-Met-Ile-Ile-Arg-Ala-Ala-Cys-His-Gly-His-Cys-Val-Pro-Asp-
 Tyr-Trp-Tyr-Glu-Thr-Ser-Phe-Trp-Leu-Leu-Trp-Ala-Asn-Ser-Ala-Val-Asn-
 Pro-Val-Leu-Tyr-Pro-Leu-Cys-His-His-Ser-Phe-Arg-Arg-Ala-Phe-Thr-Lys-
 Leu-Leu-Cys-Pro-Gln-Lys-Leu-Lys-Ile-Gln-Pro-His-Ser-Ser-Leu-Glu-Gln-
 25 Cys-Trp-Lys

The rat H3 receptor has the following sequence:

Met-Glu-Arg-Ala-Pro-Pro-Asp-Gly-Leu-Met-Asn-Ala-Ser-Gly-Thr-Leu-Ala-
 Gly-Glu-Ala-Ala-Ala-Ala-Gly-Gly-Ala-Arg-Gly-Phe-Ser-Ala-Ala-Trp-Thr-
 Ala-Val-Leu-Ala-Ala-Leu-Met-Ala-Leu-Leu-Ile-Val-Ala-Thr-Val-Leu-Gly-
 30 Asn-Ala-Leu-Val-Met-Leu-Ala-Phe-Val-Ala-Asp-Ser-Ser-Leu-Arg-Thr-Gln-
 Asn-Asn-Phe-Phe-Leu-Leu-Asn-Leu-Ala-Ile-Ser-Asp-Phe-Leu-Val-Gly-Ala-
 Phe-Cys-Ile-Pro-Leu-Tyr-Val-Pro-Tyr-Val-Leu-Thr-Gly-Arg-Trp-Thr-Phe-
 Gly-Arg-Gly-Leu-Cys-Lys-Leu-Trp-Leu-Val-Val-Asp-Tyr-Leu-Leu-Cys-Ala-
 Ser-Ser-Val-Phe-Asn-Ile-Val-Leu-Ile-Ser-Tyr-Asp-Arg-Phe-Leu-Ser-Val-
 35 Thr-Arg-Ala-Val-Ser-Tyr-Arg-Ala-Gln-Gln-Gly-Asp-Thr-Arg-Arg-Ala-Val-
 Arg-Lys-Met-Ala-Leu-Val-Trp-Val-Leu-Ala-Phe-Leu-Leu-Tyr-Gly-Pro-Ala-

Ile-Leu-Ser-Trp-Glu-Tyr-Leu-Ser-Gly-Gly-Ser-Ser-Ile-Pro-Glu-Gly-His-
 Cys-Tyr-Ala-Glu-Phe-Phe-Tyr-Asn-Trp-Tyr-Phe-Leu-Ile-Thr-Ala-Ser-Thr-
 Leu-Glu-Phe-Phe-Thr-Pro-Phe-Leu-Ser-Val-Thr-Phe-Phe-Asn-Leu-Ser-Ile-
 Tyr-Leu-Asn-Ile-Gln-Arg-Arg-Thr-Arg-Leu-Arg-Leu-Asp-Gly-Gly-Arg-Glu-
 5 Ala-Gly-Pro-Glu-Pro-Pro-Pro-Asp-Ala-Gln-Pro-Ser-Pro-Pro-Pro-Ala-Pro-
 Pro-Ser-Cys-Trp-Gly-Cys-Trp-Pro-Lys-Gly-His-Gly-Glu-Ala-Met-Pro-Leu-
 His-Arg-Tyr-Gly-Val-Gly-Glu-Ala-Gly-Pro-Gly-Val-Glu-Ala-Gly-Glu-Ala-
 Ala-Leu-Gly-Gly-Gly-Ser-Gly-Gly-Gly-Ala-Ala-Ala-Ser-Pro-Thr-Ser-Ser-
 Ser-Gly-Ser-Ser-Ser-Arg-Gly-Thr-Glu-Arg-Pro-Arg-Ser-Leu-Lys-Arg-Gly-
 10 Ser-Lys-Pro-Ser-Ala-Ser-Ser-Ala-Ser-Leu-Glu-Lys-Arg-Met-Lys-Met-Val-
 Ser-Gln-Ser-Ile-Thr-Gln-Arg-Phe-Arg-Leu-Ser-Arg-Asp-Lys-Lys-Val-Ala-
 Lys-Ser-Leu-Ala-Ile-Ile-Val-Ser-Ile-Phe-Gly-Leu-Cys-Trp-Ala-Pro-Tyr-
 Thr-Leu-Leu-Met-Ile-Ile-Arg-Ala-Ala-Cys-His-Gly-Arg-Cys-Ile-Pro-Asp-
 Tyr-Trp-Tyr-Glu-Thr-Ser-Phe-Trp-Leu-Leu-Trp-Ala-Asn-Ser-Ala-Val-Asn-
 15 Pro-Val-Leu-Tyr-Pro-Leu-Cys-His-Tyr-Ser-Phe-Arg-Arg-Ala-Phe-Thr-Lys-
 Leu-Leu-Cys-Pro-Gln-Lys-Leu-Lys-Val-Gln-Pro-His-Gly-Ser-Leu-Glu-Gln-
 Cys-Trp-Lys

The assay measures the activation of G proteins by catalyzing the exchange of
 guanosine 5'-diphosphate (GDP) by guanosine 5'-triphosphate (GTP) at the α -subunit. The
 20 GTP-bounded G proteins dissociate into two subunits, $G\alpha_{GTP}$ and $G\beta\gamma$, which in turn regulate
 intracellular enzymes and ion channels. GTP is rapidly hydrolysed by the $G\alpha$ -subunit
 (GTPases) and the G protein is deactivated and ready for a new GTP exchange cycle. To
 study the function of ligand induced G protein coupled receptor (GPCR) activation by an in-
 crease in guanine nucleotide exchange at the G proteins, the binding of [35 S]-guanosine-5'-
 25 O-(3-thio) triphosphate ([35 S] GTP γ S), a non-hydrolysed analogue of GTP, is determined.
 This process can be monitored *in vitro* by incubating cell membranes containing the G pro-
 tein coupled receptor H3 with GDP and [35 S] GTP γ S. Cell membranes are obtained from
 CHO cells stably expressing the human H3 receptor or from HEK 293 cells stably expressing
 the rat or monkey H3 receptor. The cells are washed twice in PBS, harvested with PBS+1
 30 mM EDTA, pH 7.4 and centrifuged at 280 g for 5 min. The cell pellet is homogenized in 10 ml
 ice-cold Hepes buffer (20 mM Hepes, 10 mM EDTA pH 7.4 (NaOH)) using an Ultra-Turrax
 homogenizer for 30 seconds and centrifuged for 15 min at 30.000 g. Following this centrifu-
 gation step, the membrane pellet is resuspended in 10 ml ice-cold Hepes buffer (20 mM
 Hepes, 0.1 mM EDTA pH 7.4 (NaOH)) and homogenized as describe above. This procedure
 35 is repeated twice except for the last homogenization step, the protein concentration is deter-

mined and membranes are diluted to a protein concentration of 2 mg/ml, aliquoted and kept at -80°C until use.

In order to study the presence and the potency of an inverse agonist/antagonist the H3-receptor agonist ligand R- α -methyl histamine (RAMHA) is added. The ability of the test compound to counteract the effect of RAMHA is measured. When studying the effect of an agonist RAMHA is not added to the assay medium. The test compound is diluted in the assay buffer (20 mM HEPES, 120 mM NaCl, 10 mM MgCl_2 pH 7.4 (NaOH)) at various concentrations followed by addition of 10^{-8} nM RAMHA (only in the case where an inverse agonist/antagonist is examined), 3 μM GDP, 2.5 μg membranes, 0.5 mg SPA beads and 0.1 nM [^{35}S] GTP γS and incubated for 2 hours by slightly shaking at room temperature. For the rat and monkey H3 receptor 10 μg membranes including 10 $\mu\text{g}/\text{ml}$ saponin are used. The plates are centrifuged at 420 g for 10 min and the radioactivity is measured using a Top-counter. The results are analyzed by non linear regression and the IC_{50} value is determined.

RAMHA and other H3 agonists stimulate the binding of [^{35}S] GTP γS to membranes expressing the H3 receptor. In the antagonist/inverse agonist test, the ability of increasing amounts of test compound to inhibit the increased [^{35}S] GTP γS binding by 10^{-8} M RAMHA is measured as a decrease in radioactivity signal. The IC_{50} value determined for an antagonist is the ability of this compound to inhibit the effect of 10^{-8} M RAMHA by 50%. In the agonist test, the ability of increasing amounts of test compound is measured as an increase in radioactivity signal. The EC_{50} value determined for an agonist, is the ability of this compound to increase the signal by 50% of the maximal signal that is obtained by 10^{-5} M RAMHA.

Preferably, the antagonists and agonists according to the invention have an $\text{IC}_{50}/\text{EC}_{50}$ value as determined by one or more of the assays of less than 10 μM , more preferred of less than 1 μM , and even more preferred of less than 500 nM, such as of less than 100 nM.

The open cage Schedule-fed rat model

The ability of the present compounds to reduce weight is determined using the *in vivo* open cage Schedule-fed rat model.

Sprague-Dawley (SD) male rats of an age of about 1½ to 2 months and a weight of about 200-250 g are purchased from Møllegaard Breeding and Research Centre A/S (Denmark). On arrival they are allowed some days of acclimatisation before being placed in individual open plastic cages. They are habituated to the presence of food (Altromin pelleted rat chow) in their home cage only during 7 hours in the morning from 07.30 to 14.30 all days a


week. Water is present ad libitum. As the consumption of food has stabilised after 7 to 9 days, the animals are ready for use.

Each animal is used only once to avoid carry-over effects between treatments. During the test sessions, the test compound is administered intraperitoneally or orally 30 min before the start of the sessions. One group of animals is administered the test compound at different doses and a control group of animals is given a vehicle. Food and water intake are monitored at 1, 2 and 3 hours post administration.

Any side effects may rapidly be discovered (barrel-rolling, bushy fur etc.) since the animals are kept in transparent plastic cages to enable continuous monitoring.

CLAIMS

1. A compound of the general formula (I):



5 wherein

R¹ represents

- hydrogen,
- C₁₋₈-alkyl, C₃₋₈-alkenyl or C₃₋₈-alkynyl,
which may optionally be substituted with one or more halogen atoms,
- C₃₋₇-cycloalkyl, C₃₋₇-cycloalkenyl, C₄₋₈-bicycloalkyl, C₃₋₇-cycloalkyl-C₁₋₃-alkyl or
C₃₋₇-cycloalkenyl-C₁₋₃-alkyl,
which may optionally be substituted at any position with one or more halogen atoms,

R² represents C₁₋₈-alkyl,

20 f is 0, 1 or 2,

X represents $-(\text{CH}_2)_m-(\text{Z})_n-(\text{CH}_2)_o-$,

m and o independently are 0, 1, 2, 3 or 4,

25

n is 0 or 1,

Z is $-\text{O}-$, $-\text{NH}-$, $-\text{N}(\text{CH}_3)-$, $-\text{C}(=\text{O})-$, $-\text{CH}(\text{OH})-$, $-\text{CH}(\text{O}-\text{C}_{1-6}\text{-alkyl})-$, $-\text{C}(=\text{N}-\text{OH})-$, $-\text{S}-$, $-\text{S}(=\text{O})-$,
 $-\text{S}(=\text{O})_2-$, $-\text{CH}=\text{CH}-$ or $-\text{C}\equiv\text{C}-$,

30

Y is

(a) aryl or heteroaryl, which may optionally be substituted with one or more substituents selected from

- 5 ▪ halogen, nitro, cyano, hydroxy, C₁₋₆-alkanoyl, C₁₋₆-alkylthio, C₁₋₆-alkylsulfonyl, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₃₋₈-cycloalkyl, trifluoromethyl, trifluoromethoxy, -NR³R⁴ and -O(C=O)NR³R⁴, or wherein two substituents in adjacent positions form a radical -O-(CH₂)₁₋₃-O-,
 wherein R³ and R⁴ independently are hydrogen, C₁₋₆-alkyl, C₃₋₈-cycloalkyl,
 10 C₁₋₆-alkanoyl or aryl, or R³ and R⁴ together with the nitrogen atom to which they are attached form a 4 to 7 membered, saturated or unsaturated azetidiny, pyrrolidinyl, piperidyl or azepanyl ring,
- 15 ▪ aryl, aryloxy, aryl-C₁₋₆-alkyl and aryl-C₁₋₆-alkoxy, wherein the ring moieties optionally may be substituted with one or more substituents selected from
 - o halogen, nitro, cyano, hydroxy, C₁₋₆-alkanoyl, C₁₋₆-alkylthio, C₁₋₆-alkylsulfonyl, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₃₋₈-cycloalkyl, trifluoromethyl, trifluoromethoxy, -NR⁵R⁶ and -O(C=O)NR⁵R⁶, or wherein two substituents in adjacent positions form a radical -O-(CH₂)₁₋₃-O-,
 20 wherein R⁵ and R⁶ independently are hydrogen, C₁₋₆-alkyl, C₃₋₈-cycloalkyl, C₁₋₆-alkanoyl or aryl, or R⁵ and R⁶ together with the nitrogen atom to which they are attached form a 4 to 7 membered, saturated or unsaturated azetidiny, pyrrolidinyl, piperidyl or azepanyl ring,
- 25 (b) C₃₋₈-cycloalkyl or C₅₋₈-cycloalkenyl, which may optionally be substituted with one or more substituents selected from
 - C₁₋₆-alkyl, C₁₋₆-alkoxy, C₁₋₆-alkylthio, cyano, trifluoromethyl, trifluoromethoxy and halogen,
 - 30 ▪ aryl and aryloxy, wherein the ring moieties optionally may be substituted with one or more substituents selected from
 - o halogen, nitro, cyano, hydroxy, C₁₋₆-alkanoyl, C₁₋₆-alkylthio, C₁₋₆-alkylsulfonyl, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₃₋₈-cycloalkyl, trifluoromethyl, trifluoromethoxy, -NR⁷R⁸ and -O(C=O)NR⁷R⁸, or wherein two substituents in adjacent positions form a radical -O-(CH₂)₁₋₃-O-,
 35

wherein R^7 and R^8 independently are hydrogen, C_{1-6} -alkyl, C_{3-8} -cycloalkyl, C_{1-6} -alkanoyl or aryl, or R^7 and R^8 together with the nitrogen atom to which they are attached form a 4 to 7 membered, saturated or unsaturated azetid-
inyl, pyrrolidinyl, piperidyl or azepanyl ring,

5

as well as any diastereomer or enantiomer or tautomeric form thereof including mixtures of these or a pharmaceutically acceptable salt thereof,



with the proviso that,

10


when m, n, and o are 0, R^1 must not be cyclopentyl, cyclohexyl, ethyl, or methyl,




and with the proviso that the compound must not be

15




20





or

5



2. A compound according to claim 1, wherein f is 1.

10 3. A compound according to claim 1 or 2, wherein R¹ is C₁₋₈-alkyl, C₃₋₇-cycloalkyl-C₁₋₃-alkyl or C₃₋₇-cycloalkyl, which may optionally be substituted with one or more halogen atoms.

4. A compound according to claim 3, wherein R¹ is C₁₋₈-alkyl, C₃₋₇-cycloalkyl-C₁₋₃-alkyl or C₃₋₇-cycloalkyl.

15

5. A compound according to claim 4, wherein R¹ is C₁₋₈-alkyl.

6. A compound according to claim 5, wherein R¹ is 1-ethylpropyl.

20 7. A compound according to claim 5, wherein R¹ is 2-propyl.


8. A compound according to claim 4, wherein R^1 is C_{3-7} -cycloalkyl.
9. A compound according to claim 8, wherein R^1 is cyclobutyl or cyclopentyl.
- 5 10. A compound according to any one of the preceding claims, wherein R^2 is methyl.
11. A compound according to any one of the preceding claims, wherein X is $-(CH_2)_m-C(=O)-CH_2-O-$, $-(CH_2)_m-$, a valence bond or $-(CH_2)_m-CH=CH-(CH_2)_o-$, wherein m and o are as defined in claim 1.
- 10 12. A compound according to claim 11, wherein X is $-(CH_2)_m-C(=O)-$, $-(CH_2)_m-$, a valence bond or $-CH=CH-$, wherein m is 1, 2, 3 or 4.
13. A compound according to claim 11, wherein X is $-(CH_2)_m-C(=O)-$, $-(CH_2)_m-$ or $-CH=CH-$,
15 wherein m is 1, 2, 3 or 4.
14. A compound according to claim 11, wherein X is $-(CH_2)_2-C(=O)-$, $-CH_2-$, $-(CH_2)_2-$, a valence bond or $-CH=CH-$.
- 20 15. A compound according to claim 11, wherein X is $-(CH_2)_2-C(=O)-$, $-CH_2-$, $-(CH_2)_2-$ or $-CH=CH-$.
16. A compound according to any one of the preceding claims, wherein Y is aryl, which may optionally be substituted as defined in claim 1.
- 25 17. A compound according to claim 16, wherein Y is phenyl, which may optionally be substituted as defined in claim 1.
18. A compound according to claim 17, wherein Y is phenyl, which may optionally be substituted with one or two of
- 30
- C_{1-6} -alkoxy, $-NR^3R^4$, halogen, trifluoromethyl or hydroxy, wherein R^3 and R^4 are as defined in claim 1,
 - two substituents in adjacent positions forming a radical $-O-(CH_2)_{1-3}-O-$, or
- 35

- phenyl or phenoxy, which may optionally be substituted as defined in claim 1.

19. A compound according to claim 18, wherein Y is phenyl, which may optionally be substituted with one or two of


- C₁₋₆-alkoxy, -NR³R⁴, halogen, trifluoromethyl or hydroxy, wherein R³ and R⁴ independently are hydrogen or C₁₋₆-alkyl,
- two substituents in adjacent positions forming a radical -O-CH₂-O-, or
- phenyl or phenoxy.

20. A compound according to claim 1 of the general formula (I₁):




wherein R¹, R² and Y are as defined in any one of the preceding claims.

21. A compound according to claim 1 of the general formula (I₂):



22. A compound according to claim 1 of the general formula (I₃):




23. Use of a compound according to any one of the preceding claims 1 to 22 as a pharmaceutical composition.

24. A pharmaceutical composition comprising, as an active ingredient, at least one compound according to any one of the claims 1 to 22 together with one or more pharmaceutically acceptable carriers or excipients.

25. A pharmaceutical composition according to claim 24 in unit dosage form, comprising from about 0.05 mg to about 1000 mg, preferably from about 0.1 mg to about 500 mg and especially preferred from about 0.5 mg to about 200 mg of the compound according to any one of the claims 1 to 21.

26. Use of a compound of the general formula (I'':



wherein

R¹ represents

- hydrogen,
- C₁₋₈-alkyl, C₃₋₈-alkenyl or C₃₋₈-alkynyl, which may optionally be substituted with one or more halogen atoms,

- C₃₋₇-cycloalkyl, C₃₋₇-cycloalkenyl, C₄₋₈-bicycloalkyl, C₃₋₇-cycloalkyl-C₁₋₃-alkyl or C₃₋₇-cycloalkenyl-C₁₋₃-alkyl, which may optionally be substituted at any position with one or more halogen atoms,

5 R² represents C₁₋₆-alkyl,

X represents $-(CH_2)_m-(Z)_n-(CH_2)_o-$,

m and o independently are 0, 1, 2, 3 or 4,

10

n is 0 or 1,

Z is -O-, -NH-, -N(CH₃)-, -C(=O)-, -CH(OH)-, -C(=N-OH)-, -S-, -S(=O)-, -S(=O)₂-, -CH=CH- or -C≡C-,

15

Y is

(a) aryl or heteroaryl, which may optionally be substituted with one or more substituents selected from

- 20
- halogen, nitro, cyano, hydroxy, C₁₋₆-alkanoyl, C₁₋₆-alkylthio, C₁₋₆-alkylsulfonyl, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₃₋₈-cycloalkyl, trifluoromethyl, trifluoromethoxy, -NR³R⁴ and -O(C=O)NR³R⁴, or wherein two substituents in adjacent positions form a radical -O-(CH₂)₁₋₃-O-, wherein R³ and R⁴ independently are hydrogen, C₁₋₆-alkyl, C₃₋₈-cycloalkyl,
- 25
- C₁₋₆-alkanoyl or aryl, or R³ and R⁴ together with the nitrogen atom to which they are attached form a 4 to 7 membered, saturated or unsaturated azetidiny, pyrrolidinyl, piperidyl or azepanyl ring,
- 30
- aryl, aryloxy, aryl-C₁₋₆-alkyl and aryl-C₁₋₆-alkoxy, wherein the ring moieties optionally may be substituted with one or more substituents selected from
 - o halogen, nitro, cyano, hydroxy, C₁₋₆-alkanoyl, C₁₋₆-alkylthio, C₁₋₆-alkylsulfonyl, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₃₋₈-cycloalkyl, trifluoromethyl, trifluoromethoxy, -NR⁵R⁶ and -O(C=O)NR⁵R⁶, or wherein two substituents in adjacent positions form a radical -O-(CH₂)₁₋₃-O-,

wherein R⁵ and R⁶ independently are hydrogen, C₁₋₆-alkyl, C₃₋₈-cycloalkyl, C₁₋₆-alkanoyl or aryl, or R⁵ and R⁶ together with the nitrogen atom to which they are attached form a 4 to 7 membered, saturated or unsaturated azetidiny, pyrrolidinyl, piperidyl or azepanyl ring,

5

(b) C₃₋₈-cycloalkyl or C₅₋₈-cycloalkenyl, which may optionally be substituted with one or more substituents selected from

- C₁₋₆-alkyl, C₁₋₆-alkoxy, C₁₋₆-alkylthio, cyano, trifluoromethyl, trifluoromethoxy and halogen,

10

- aryl and aryloxy, wherein the ring moieties optionally may be substituted with one or more substituents selected from

- halogen, nitro, cyano, hydroxy, C₁₋₆-alkanoyl, C₁₋₆-alkylthio, C₁₋₆-alkylsulfonyl, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₃₋₈-cycloalkyl, trifluoromethyl, trifluoromethoxy, -NR⁷R⁸ and -O(C=O)NR⁷R⁸, or wherein two substituents in adjacent positions form a radical -O-(CH₂)₁₋₃-O-,

15

wherein R⁷ and R⁸ independently are hydrogen, C₁₋₆-alkyl, C₃₋₈-cycloalkyl, C₁₋₆-alkanoyl or aryl, or R⁷ and R⁸ together with the nitrogen atom to which they are attached form a 4 to 7 membered, saturated or unsaturated azetidiny, pyrrolidinyl, piperidyl or azepanyl ring,

20

a well as any diastereomer or enantiomer or tautomeric form thereof including mixtures of these or a pharmaceutically acceptable salt thereof for the preparation of a pharmaceutical composition for the treatment of disorders and diseases related to the histamine H₃ receptor.

25

27. Use of a compound as defined in claim 26 for the preparation of a pharmaceutical composition for the treatment of diseases and disorders in which an inhibition of the H₃ histamine receptor has a beneficial effect.

30

28. Use of a compound as defined in claim 26 for the preparation of a pharmaceutical composition having histamine H₃ antagonistic activity or histamine H₃ inverse agonistic activity.

29. Use of a compound as defined in claim 26 for the preparation of a pharmaceutical composition for the reduction of weight.

35

30. Use of a compound as defined in claim 26 for the preparation of a pharmaceutical composition for the treatment of overweight or obesity.
- 5 31. Use of a compound as defined in claim 26 for the preparation of a pharmaceutical composition for the suppression of appetite or for satiety induction.
32. Use of a compound as defined in claim 26 for the preparation of a pharmaceutical composition for the prevention and/or treatment of disorders and diseases related to overweight or obesity.
- 10 33. Use of a compound as defined in claim 26 for the preparation of a pharmaceutical composition for the prevention and/or treatment of eating disorders such as bulimia and binge eating.
- 15 34. Use of a compound as defined in claim 26 for the preparation of a pharmaceutical composition for the treatment of IGT.
35. Use of a compound as defined in claim 26 for the preparation of a pharmaceutical composition for the treatment of type 2 diabetes.
- 20 36. Use of a compound as defined in claim 26 for the preparation of a pharmaceutical composition for the delaying or prevention of the progression from IGT to type 2 diabetes.
- 25 37. Use of a compound as defined in claim 26 for the preparation of a pharmaceutical composition for the delaying or prevention of the progression from non-insulin requiring type 2 diabetes to insulin requiring type 2 diabetes.
- 30 38. Use of a compound as defined in claim 26 for the preparation of a pharmaceutical composition for the treatment of diseases and disorders in which a stimulation of the H3 histamine receptor has a beneficial effect.
39. Use of a compound as defined in claim 26 for the preparation of a pharmaceutical composition having histamine H3 agonistic activity.

40. Use of a compound as defined in claim 26 for the preparation of a pharmaceutical composition for the treatment of allergic rhinitis, ulcer or anorexia.

5 41. Use of a compound as defined in claim 26 for the preparation of a pharmaceutical composition for the treatment of Alzheimer's disease, narcolepsy or attention deficit disorder.

42. A method for the treatment of disorders or diseases related to the H3 histamine receptor the method comprising administering to a subject in need thereof an effective amount of a compound as defined in claim 26 or a pharmaceutical composition according to claim 24 or
10 25.

43. The method according to claim 42 wherein the effective amount of the compound is in the range of from about 0.05 mg to about 2000 mg, preferably from about 0.1 mg to about 1000 mg and especially preferred from about 0.5 mg to about 500 mg per day.

15

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
27 March 2003 (27.03.2003)

PCT

(10) International Publication Number
WO 2003/024928 A3

(51) International Patent Classification⁷: **C07D 207/14**,
211/56, 405/12, 205/04, A61K 31/40, 31/445, 31/4523,
A61P 1/04, 3/04, 3/10, 25/28, 37/08

(21) International Application Number:
PCT/DK2002/000593

(22) International Filing Date:
11 September 2002 (11.09.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PA 2001 01344 14 September 2001 (14.09.2001) DK
PA 2002 00750 16 May 2002 (16.05.2002) DK

(71) Applicants: **NOVO NORDISK A/S** [DK/DK]; Novo
Allé, DK-2880 Bagsværd (DK). **BOEHRINGER IN-
GELHEIM INTERNATIONAL GMBH** [DE/DE];
55216 Ingelheim am Rhein (DE).

(72) Inventors: **DÖRWALD, Florencio, Zaragoza**; Højager-
parken 30, 1, DK-2750 Ballerup (DK). **HOHLWEG, Rolf**;
Nybovej 6, DK-3490 Kvistgaard (DK).

(74) Common Representative: **NOVO NORDISK A/S**; Cor-
porate Patents, Novo Allé, DK-2880 Bagsværd (DK).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC,
VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).


Published:

— with international search report

(88) Date of publication of the international search report:
4 March 2004

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: NOVEL AMINOAZETIDINE, -PYRROLIDINE AND -PIPERIDINE DERIVATIVES



H3 receptor.

(57) Abstract: Novel aminoazetidine, -pyrrolidine, and -piperidine derivatives of the general formula (I): use of these compounds as pharmaceutical compositions, pharmaceutical compositions comprising the compounds, and a method of treatment employing these compounds and compositions. The compounds show a high and selective binding affinity to the histamine H3 receptor indicating histamine H3 receptor antagonistic, inverse agonistic or agonistic activity. As a result, the compounds are useful for the treatment of diseases and disorders related to the histamine

INTERNATIONAL SEARCH REPORT

International Application No
PCT/DK 02/00593

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C07D207/14 C07D211/56 C07D405/12 C07D205/04 A61K31/40 A61K31/445 A61K31/4523 A61P1/04 A61P3/04 A61P3/10 A61P25/28 A61P37/08		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07D A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, WPI Data, PAJ, CHEM ABS Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 02 22592 A (SCHERING CORP) 21 March 2002 (2002-03-21) page 56, example 63, page 3, line 30 - page 4, line 3, the claims ---	1-43
P,X	WO 02 49648 A (SCHERING CORP) 27 June 2002 (2002-06-27) page 11, line 20 - page 12, line 2; the claims; the examples ---	1-43
P,A	WO 02 06223 A (ABBOTT LAB) 24 January 2002 (2002-01-24) the whole document ---	1-43
X	US 4 049 821 A (FUNDERBURK JR WILLIAM HENRY) 20 September 1977 (1977-09-20) column 1, line 15 - line 21, Table I, example 9; the claims ---	1-25,40
-/--		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "Z" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
5 December 2002		24 02 2003
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3018		Authorized officer GERD STRANDELL/ ELY

INTERNATIONAL SEARCH REPORT

Inte.al Application No

PCT/DK 02/00593

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DE 19 64 516 A (ROBINS CO INC A H) 9 July 1970 (1970-07-09) page 2; the examples; the claims ---	1-25
X	US 4 109 005 A (ALPHIN REEVIS S ET AL) 22 August 1978 (1978-08-22) the whole document ---	1-25
X	DATABASE STN INTERNATIONAL [Online] File CAPLUS, CAPLUS accession number 1978:32120, document number 88:32120; GRAYBIEL, ASHTON ET AL: "Evaluation of a new antinauseant drug for the prevention of motion sickness" XP002223803 & Aviat., Space Environ. Med. (1977), 48(9), 857-71 abstract ---	1-25
X	GB 1 364 231 A (ROBINS CO INC A H) 21 August 1974 (1974-08-21) the examples; the claims ---	1-25
X	DE 25 58 348 A (ROBINS CO INC A H) 8 July 1976 (1976-07-08) the whole document ---	1-25
X	GB 1 088 531 A (ILE DE FRANCE) 25 October 1967 (1967-10-25) the whole document -----	1-25

INTERNATIONAL SEARCH REPORT

International application No.
PCT/DK 02/00593

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: **23, 42, 43**
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Claims Nos.: 23,42,43

Claims 23,42,43 relate to methods of treatment of the human or animal body by surgery or by therapy/diagnostic methods practised on the human or animal body/Rule 39.1.(iv). Nevertheless, a search has been executed for these claims. The search has been based on the alleged effects of the compounds/composition.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/DK 02/00593

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0222592	A	21-03-2002	AU 9454701 A WO 0222592 A2 US 2002165223 A1	26-03-2002 21-03-2002 07-11-2002
WO 0249648	A	27-06-2002	AU 3405602 A WO 0249648 A1	01-07-2002 27-06-2002
WO 0206223	A	24-01-2002	AU 7338401 A WO 0206223 A1	30-01-2002 24-01-2002
US 4049821	A	20-09-1977	AU 516329 B2 AU 2510777 A BE 854507 A1 DE 2721259 A1 DK 207677 A ES 458709 A1 FI 771499 A FR 2421613 A1 GB 1582163 A IE 45384 B1 JP 52142061 A NL 7705215 A NZ 184073 A PH 13688 A PT 66532 A ,B SE 7705445 A ZA 7702527 A	28-05-1981 16-11-1978 01-09-1977 24-11-1977 13-11-1977 01-03-1978 13-11-1977 02-11-1979 31-12-1980 11-08-1982 26-11-1977 15-11-1977 20-03-1985 01-09-1980 01-06-1977 13-11-1977 26-04-1978
DE 1964516	A	09-07-1970	CH 532571 A DE 1964516 A1 FR 2026924 A5 FR 2182929 A2 FR 2182930 A2 GB 1364231 A GB 1281188 A JP 49031985 B US 3577440 A US RE29828 E	15-01-1973 09-07-1970 25-09-1970 14-12-1973 14-12-1973 21-08-1974 12-07-1972 27-08-1974 04-05-1971 07-11-1978
US 4109005	A	22-08-1978	AU 518915 B2 AU 3204877 A BE 861790 A1 JP 1437006 C JP 53139728 A JP 62041205 B PH 12742 A PT 66911 A ,B ZA 7704212 A	29-10-1981 05-07-1979 31-03-1978 25-04-1988 06-12-1978 02-09-1987 09-08-1979 01-09-1977 28-06-1978
GB 1364231	A	21-08-1974	CA 999003 A1 CA 998398 A2 CH 580582 A5 CH 590227 A5 DE 2316644 A1 DE 2365904 A1 DK 138491 B ES 413260 A1 FR 2182929 A2	26-10-1976 12-10-1976 15-10-1976 29-07-1977 18-10-1973 12-05-1977 18-09-1978 16-05-1976 14-12-1973

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/DK 02/00593

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
GB 1364231	A		FR 2182930 A2	14-12-1973
			JP 1041154 C	23-04-1981
			JP 50069066 A	09-06-1975
			JP 55027905 B	24-07-1980
			JP 50069067 A	09-06-1975
			JP 1094725 C	27-04-1982
			JP 49013165 A	05-02-1974
			JP 56020312 B	13-05-1981
			JP 51082261 A	19-07-1976
			JP 1218623 C	17-07-1984
			JP 56127354 A	06-10-1981
			JP 58046512 B	17-10-1983
			NL 7304557 A ,B,	05-10-1973
			NO 142077 B	17-03-1980
			PH 15259 A	02-11-1982
			SE 402102 B	19-06-1978
			SE 431868 B	05-03-1984
			SE 7711953 A	24-10-1977
			SE 431869 B	12-03-1984
			SE 8302193 A	19-04-1983
			US 4282243 A	04-08-1981
			US 4282244 A	04-08-1981
			US 3966957 A	29-06-1976
			US 3963745 A	15-06-1976
			US 4279822 A	21-07-1981
			ZA 7302296 A	27-11-1974
			CH 532571 A	15-01-1973
			DE 1964516 A1	09-07-1970
			FR 2026924 A5	25-09-1970
			GB 1281188 A	12-07-1972
			JP 49031985 B	27-08-1974
			US 3577440 A	04-05-1971
			US RE29828 E	07-11-1978

DE 2558348	A	08-07-1976	US 4002757 A	11-01-1977
			AU 499075 B2	05-04-1979
			AU 8755075 A	23-06-1977
			BE 836975 A1	16-04-1976
			CA 1071636 A1	12-02-1980
			CH 611603 A5	15-06-1979
			DE 2558348 A1	08-07-1976
			DK 589075 A ,B,	27-06-1976
			ES 443845 A1	16-04-1977
			FI 753606 A ,B,	27-06-1976
			FR 2295744 A1	23-07-1976
			GB 1537848 A	04-01-1979
			IE 43819 B1	03-06-1981
			JP 1345893 C	29-10-1986
			JP 60248684 A	09-12-1985
			JP 61009314 B	22-03-1986
			JP 1336353 C	11-09-1986
			JP 51091254 A	10-08-1976
			JP 61001426 B	17-01-1986
			NL 7515084 A	29-06-1976
			NO 754376 A ,B,	29-06-1976
			PH 13883 A	27-10-1980
			SE 408417 B	11-06-1979
			SE 7514374 A	28-06-1976

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/DK 02/00593

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 2558348	A	US 4039678 A	02-08-1977
		US 4087428 A	02-05-1978
		US 4073796 A	14-02-1978
		ZA 7507972 A	29-12-1976

GB 1088531	A	25-10-1967	US 3342826 A
			19-09-1967
			BE 658250 A
			13-07-1965
			CH 468377 A
			15-02-1969
			DE 1795723 A1
			29-08-1974
			DE 1595915 A1
			12-02-1970
			DK 124402 B
			16-10-1972
			FI 48273 B
			30-04-1974
			FR 4879 M
			FR 1472025 A
			24-05-1967
			IL 22669 A
			27-03-1969
			NL 6500326 A
			14-07-1965
			NO 119533 B
			01-06-1970
			QA 1859 A
			14-01-1970
			SE 340452 B
			22-11-1971
